

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/51</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 97/07135</b> <b>(43) International Publication Date:</b> 27 February 1997 (27.02.97)
<b>(21) International Application Number:</b> PCT/US96/13163 <b>(22) International Filing Date:</b> 13 August 1996 (13.08.96) <b>(30) Priority Data:</b> 60/002,313 14 August 1995 (14.08.95) US <b>(71) Applicants:</b> CREATIVE BIOMOLECULES, INC. [US/US]; 45 South Street, Hopkinton, MA 01748 (US). THE LUD- WIG INSTITUTE FOR CANCER RESEARCH [US/US]; 1345 Avenue of the Americas, New York, NY 10105 (US). <b>(72) Inventors:</b> ICHJO, Hidenori; The Cancer Institute, Tokyo, Japanese Foundation for Cancer Research, Dept. of Bio- chemistry, 1-37-1 Kami-ikebukuro, Toshima-ku, Tokyo 170 (JP). NISHITOH, Hideki; The Cancer Institute, Tokyo, Japanese Foundation for Cancer Research, Dept. of Bio- chemistry, 1-37-1 Kami-ikebukuro, Toshima-ku, Tokyo 170 (JP). SAMPATH, Kuber, T.; 6 Spring Street, Medway, MA 02053 (US). <b>(74) Agent:</b> GREENHALGH, Duncan, A.; Testa, Hurwitz & Thibault, L.L.P., High Street Tower, 125 High Street, Boston, MA 02110 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> BINDING OF OSTEOGENIC PROTEIN-I (OP-1) AND ANALOGS THEREOF TO THE CELL SURFACE RECEPTOR ALK-1 AND ANALOGS THEREOF  <b>(57) Abstract</b>  Disclosed are (1) nucleic acid sequences, amino acid sequences, homologies, structural features, and various other data characterizing morphogen cell surface receptors, particularly OP-1-binding cell surface receptors; (2) methods for producing receptor proteins, including fragments thereof, using recombinant DNA technology; (3) methods for identifying novel morphogen receptors and their encoding DNAs; (4) methods for identifying compounds capable of modulating endogenous morphogen receptor levels; and (5) methods and compositions for identifying and producing morphogen analogs useful in the design of morphogen agonists and antagonists for therapeutic, diagnostic, and experimental uses.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**BINDING OF OSTEOGENIC PROTEIN-1 (OP-1) AND ANALOGS THEREOF TO THE CELL SURFACE RECEPTOR ALK-1 AND ANALOGS THEREOF**

**Field of the Invention**

This invention relates generally to the field of tissue morphogenesis and more particularly to morphogenic protein-specific cell surface receptors.

**5 Related Applications**

The present invention claims priority to Provisional Application U.S.S.N. 60/002,313 filed August 14, 1995, the entire disclosure of which is incorporated herein by reference.

**Background of the Invention**

10 Cell differentiation is the central characteristic of tissue morphogenesis which initiates during embryogenesis, and continues to various degrees throughout the life of an organism in adult tissue repair and regeneration mechanisms. The degree of morphogenesis in adult tissue varies among different tissues and is related, among other things, to the degree of cell turnover in a given tissue.

15 The cellular and molecular events which govern the stimulus for differentiation of cells is an area of intensive research. In the medical and veterinary fields, it is anticipated that the discovery of the factor or factors which control cell differentiation and tissue morphogenesis will advance significantly medicine's ability to repair and regenerate diseased or damaged mammalian tissues and organs. Particularly useful areas for human and veterinary therapeutics include reconstructive surgery and in the treatment of tissue degenerative diseases including arthritis,  
20 emphysema, osteoporosis, cardiomyopathy, cirrhosis, degenerative nerve diseases, inflammatory diseases, and cancer, and in the regeneration of tissues, organs and limbs. (In this and related applications, the terms "morphogenetic" and "morphogenic" are used interchangeably.)

A number of different factors have been isolated in recent years which appear to play a role in cell differentiation. Recently, a distinct subfamily of the "superfamily" of structurally

- 2 -

related proteins referred to in the art as the "transforming growth factor- $\beta$  (TGF- $\beta$ ) superfamily of proteins" have been identified as true tissue morphogens.

The members of this distinct "subfamily" of true tissue morphogenic proteins share substantial amino acid sequence homology within their morphogenically active C-terminal domains (at least 50% identity in the C-terminal 102 amino acid sequence), including a conserved six or seven cysteine skeleton, and share the *in vivo* activity of inducing tissue-specific morphogenesis in a variety of organs and tissues. The proteins apparently contact and interact with progenitor cells *e.g.*, by binding suitable cell surface molecules, predisposing or otherwise stimulating the cells to proliferate and differentiate in a morphogenically permissive environment. These morphogenic proteins are capable of inducing the developmental cascade of cellular and molecular events that culminate in the formation of new organ-specific tissue, including any vascularization, connective tissue formation, and nerve innervation as required by the naturally occurring tissue. The proteins have been shown to induce morphogenesis of both bone cartilage and bone, as well as periodontal tissues, dentin, liver, and neural tissue, including retinal tissue.

The true tissue morphogenic proteins identified to date include proteins originally identified as bone inductive proteins. These include OP-1, (osteogenic protein-1, also referred to in related applications as "OP1"), its *Drosophila* homolog, 60A, with which it shares 69% identity in the C-terminal "seven cysteine" domain, and the related proteins OP-2 (also referred to in related applications as "OP2") and OP-3, both of which share approximately 70-75% identity with OP-1 in the C-terminal seven cysteine domain, as well as BMP-5, BMP-6 and its murine homolog, Vgr-1, all of which share greater than 85% identity with OP-1 in the C-terminal seven cysteine domain, and the BMP-6 *Xenopus* homolog, Vgl, which shares approximately 57% identity with OP-1 in the C-terminal seven cysteine domain. Other bone inductive proteins include the CBMP-2 proteins (also referred to in the art as BMP-2 and BMP-4) and their *Drosophila* homolog, dpp. Another tissue morphogenic protein is GDF-1 (from mouse). See, for example, PCT documents US92/01968 and US92/07358, the disclosures of which are incorporated herein by reference. Other proteins identified in the art as bone morphogenic proteins include BMP-9 (See PCT/US92/05374; WO 93/00432); BMP-10 (PCT/US94/05288; WO 94/26893); BMP-11 (PCT/US94/05290; WO 94/26892); and BMP-12 and BMP-13 (PCT/US94/14030; WO 95/16035).

- 3 -

As stated above, these true tissue morphogenic proteins are recognized in the art as a distinct subfamily of proteins different from other members of the TGF- $\beta$  superfamily in that they share a high degree of sequence identity in the C-terminal domain and in that the true tissue morphogenic proteins are able to induce, on their own, the full cascade of events that result in formation of functional tissue rather than merely inducing formation of fibrotic (scar) tissue. Specifically, members of the family of morphogenic proteins are capable of all of the following in a morphogenically permissive environment: stimulating cell proliferation and cell differentiation, and supporting the growth and maintenance of differentiated cells. The morphogenic proteins apparently may act as endocrine, paracrine or autocrine factors.

The morphogenic proteins are capable of significant species "crosstalk." That is, xenogenic (foreign species) homologs of these proteins can substitute for one another in functional activity. For example, dpp and 60A, two *Drosophila* proteins, can substitute for their mammalian homologs, BMP-2/4 and OP-1, respectively, and induce endochondral bone formation at a non-bony site in a standard rat bone formation assay. Similarly, BMP-2 has been shown to rescue a *dpp*<sup>-</sup> mutation in *Drosophila*. In their native form, however, the proteins appear to be tissue-specific, each protein typically being expressed in or provided to one or only a few tissues or, alternatively, expressed only at particular times during development. For example, GDF-1 appears to be expressed primarily in neural tissue, while OP-2 appears to be expressed at relatively high levels in early (*e.g.*, 8-day) mouse embryos. The endogenous morphogens may be synthesized by the cells on which they act, by neighboring cells, or by cells of a distant tissue, the secreted protein being transported to the cells to be acted on.

A particularly potent tissue morphogenic protein is OP-1. This protein, and its xenogenic homologs, are expressed in a number of tissues, primarily in tissues of urogenital origin, as well as in bone, mammary and salivary gland tissue, reproductive tissues, and gastrointestinal tract tissue. It is also expressed in different tissues during embryogenesis, its presence coincident with the onset of morphogenesis of that tissue.

The morphogenic protein signal transduction across a cell membrane appears to occur as a result of specific binding interaction with one or more cell surface receptors. Recent studies on cell surface receptor binding of various members of the TGF- $\beta$  protein superfamily suggests that the ligands can mediate their activity by interaction with two different receptors, referred to as

- 4 -

Type I and Type II receptors to form a hetero-complex. A cell surface bound beta-glycan also may enhance the binding interaction. The Type I and Type II receptors are both serine /threonine kinases, and share similar structures: an intracellular domain that consists essentially of the kinase, a short, extended hydrophobic sequence sufficient to span the membrane one time, and an  
5 extracellular domain characterized by a high concentration of conserved cysteines.

A number of Type II receptor sequences recently have been identified. These include "TGF- $\beta$ R II", a TGF- $\beta$  Type II receptor (Lin *et al.* (1992) *Cell* 68: 775-785); and numerous activin-binding receptors. See, for example, Mathews *et al.* (1991) *Cell* 65: 973-982 and international patent application WO 92/20793, published November 26, 1992, disclosing the  
10 "ActR II" sequence; Attisano *et al.*, (1992) *Cell* 68: 97-108, disclosing the "ActR-IIB" sequence; and Legerski *et al.* (1992) *Biochem Biophys. Res. Commu.* 183: 672-679. A different Type II receptor shown to have affinity for activin is Atr-II (Childs *et al.* (1993) *PNAS* 90: 9475-9479.) Two Type II receptors have been identified in *C. elegans*, the *daf-1* gene, (Georgi *et al.* (1990) *Cell* 61: 635-645), having no known ligand to date, and *daf-4*, which has been shown to bind  
15 BMP-4, but not activin or TGF- $\beta$  (Estevez, *et al.* (1993) *Nature* 365: 644-649.)

Ten Dijke *et al.* disclose the cloning of six different Type I cell surface receptors from murine and human cDNA libraries ((1993) *Oncogene* 8: 2879-2887, and (1994) *Science* 264: 101-104). These receptors, referenced as ALK-1 to ALK-6 ("activin receptor-like kinases"), share significant sequence identities (60-79%) and several have been identified as TGF- $\beta$  binding  
20 (ALK-5) or activin binding (ALK-2, ALK-4) receptors. Xie *et al.* also report a *Drosophila* Type I receptor encoded by the *sax* gene ((1994) *Science* 263: 1756-1759). The authors suggest that the protein binds dpp.

To date, a ligand having binding affinity for the ALK-1 Type I receptor has not yet been identified. It is an object of this invention to provide molecules, including OP-1 and OP-1-related  
25 proteins, which are competent to act as ligands for ALK-1 receptor binding and/or are capable of mediating an intracellular effect by interaction with the ALK-1 receptor. Another object is to provide methods for identifying genes and naturally occurring sequences in a variety of species and/or tissues, and in a variety of nucleic acid libraries encoding ALK-1-binding ligands and ALK-1-related receptors. Still another object is to provide means for designing biosynthetic receptor-  
30 binding ligand analogs, particularly OP-1 and OP-1-related analogs, and/or for identifying natural-

- 5 -

occurring ligand analogs, including agonists and antagonists, using the receptor molecules described herein, and analogs thereof. Another object is to provide antagonists, including soluble receptor constructs comprising the extracellular ligand-binding domain, which can modulate the availability of OP-1 or an OP-1-related protein for ALK-1 receptor binding *in vivo*. Yet another

5 object is to provide means and compositions for ligand affinity purification and for diagnostic detection and quantification of ligands in a body fluid using an ALK-1 receptor and ligand-binding fragments thereof. Still another object is to provides means and compositions for modulating the endogenous expression or concentration of these receptor molecules. Yet another object is to

10 provide ligand-receptor complexes and analog sequences thereof, as well as antibodies capable of identifying and distinguishing the complex from its component proteins. Still another object is to provide means and compositions for modulating a morphogenesis in a mammal. These and other objects and features of the invention will be apparent from the description, drawings and claims which follow.

### Summary of the Invention

15 It now has been discovered that the Type I cell surface receptor molecule identified in the literature as ALK-1 is capable of specific binding affinity with true tissue morphogenic proteins, particularly OP-1 and OP-1-related proteins. Accordingly, the invention provides ligand-receptor complexes comprising at least the ligand binding domain of an ALK-1 receptor and OP-1 or an

20 OP-1 related protein or analog thereof as the ligand. The invention also provides means for identifying and/or designing useful receptor sequence analogs and OP-1, OP-1-related proteins, and analogs hereof, and means for modulating the tissue morphogenesis capability of a cell.

The ALK-1 receptor (Seq. ID No. 1) useful in this invention is part of a family of receptor molecules which share a conserved structure, including an extracellular, ligand binding domain generally composed of about 100-130 amino acids (Type I receptors; up to about 196 amino acids

25 for Type II receptors), a transmembrane domain sufficient to span a cellular membrane one time, and an intracellular (cytoplasmic) domain having serine/threonine kinase activity. The intact receptor is a single polypeptide chain of about 500-550 amino acids and having an apparent molecular weight of about 50-55 kDa.

As used herein, ligand-receptor binding specificity is understood to mean a specific,

30 saturable noncovalent interaction between the ligand and the receptor, and which is subject to

competitive inhibition by a suitable competitor molecule. Preferred binding affinities (defined as the amount of ligand required to fill one-half (50%) of available receptor binding sites) are described herein by dissociation constant ( $K_d$ ). In one embodiment, preferred binding affinities of the ligand-receptor complexes described herein have a  $K_d$  of less than  $10^{-7}$  M, preferably less than  $5 \times 10^{-7}$  M, more preferably less than  $10^{-8}$  M. In another preferred embodiment, the receptor molecules have little or no substantial binding affinity for TGF- $\beta$ .

As will be appreciated by those having ordinary skill in the art, the ALK-1 receptor can also have binding affinity for other, related morphogenic proteins. It further is contemplated that one or more related morphogenic proteins may have better binding affinity and/or signal transducing ability than OP-1 has for ALK-1. As used herein, an OP-1-related protein/ALK-1 ligand is understood to have substantially the same binding affinity for ALK-1 as OP-1 if it can be competed successfully for ALK-1 binding in a standard competition assay with OP-1. In one preferred embodiment, the ligand is OP-1 or an OP-1-related protein or analog and has a binding affinity for ALK-1 defined by a dissociation constant of less than about  $10^{-7}$  M, preferably less than about  $5 \times 10^{-7}$  M or  $10^{-8}$  M. It is anticipated however, that analogs having lower binding affinities, e.g., on the order of  $10^{-6}$  M also will be useful. For example, such analogs may be provided to a mammal to modulate availability of serum-soluble ligand for ALK-1 receptor binding *in vivo*. Similarly, where tight binding interaction is desired, for example as part of a cancer therapy wherein the analog acts as a ligand-receptor antagonist, preferred binding affinities may be on the order of  $5 \times 10^{-8}$  M.

In another embodiment, the ALK-1 receptor binding analogs contemplated by the invention include proteins encoded by nucleic acids which hybridize with the DNA sequence encoding OP-1 under stringent hybridization conditions, and which have substantially the same or better binding affinity for a ALK-1 as OP-1. As used herein, stringent hybridization conditions are as defined in the art, (see, for example, *Molecular Cloning: A Laboratory Manual*, Maniatis *et al.*, eds. 2d. ed., Cold Spring Harbor Press, Cold Spring Harbor, 1989.) An exemplary set of conditions is defined as: hybridization in 40% formamide, 5xSSPE, 5xDenhardt's Solution, and 0.1% SDS at 37°C overnight, and washing in 0.1xSSPE, 0.1% SDS at 50°C.

Useful receptor analogs include xenogenic (foreign species) homologs of the human ALK-1 sequences described herein, including those obtained from other mammalian species, as well as



- 7 -

other, eukaryotic, non-mammalian xenogenic homologs. Also contemplated are biosynthetic constructs and naturally-occurring sequence variants of ALK-1, provided these molecules bind OP-1 or an OP-1 related protein, or analog specifically as defined herein. In one embodiment, sequence variants include receptor analogs which have at least the same binding affinity for OP-1 as ALK-1 and which are recognized by an antibody having binding specificity for ALK-1.

The ALK-1 receptor analogs may be used as an OP-1 or an OP-1-related protein antagonist. For example, a soluble form of a receptor, *e.g.*, consisting essentially of only the extracellular ligand binding domain, may be provided systemically to a mammal to bind to soluble ligand, effectively competing with ligand binding to a cell surface receptor, thereby modulating (reducing) the availability of free ligand *in vivo* for cell surface binding.

The true tissue morphogenic proteins contemplated as useful receptor ligands in the invention include OP-1, OP-1-related proteins, and analogs thereof. As used herein, an "analog" is understood to include all molecules able to functionally substitute for OP-1 or an OP-1-related protein in Type I receptor binding, *e.g.*, which are able to compete successfully with OP-1 for receptor binding in a standard competition assay. In one embodiment, useful analogs include molecules whose binding affinity is defined by a dissociation constant of less than about  $5 \times 10^{-6}$  M, preferably less than about  $10^{-7}$  M or  $5 \times 10^{-7}$  M. As described above, both stronger and weaker binding affinities are contemplated to be useful in particular applications.

The OP-1 analogs contemplated herein, all of which mimic the binding activity of OP-1 or an OP-1-related protein sufficiently to act as a substitute for OP-1 in receptor binding, can act as OP-1 agonists, capable of mimicking OP-1 both in receptor binding and in inducing a transmembrane effect *e.g.*, inducing threonine- or serine-specific phosphorylation following binding. Alternatively, the OP-1 analog can act as an OP-1 antagonist, capable of mimicking OP-1 in receptor binding only, but unable to induce a transmembrane effect, thereby blocking the natural ligand from interacting with its receptor, for example. Useful applications for antagonists include their use as therapeutics to modulate uncontrolled differentiated tissue growth, such as occurs in malignant transformations such as in osteosarcomas or Paget's disease.

OP-1 analogs contemplated by the invention can be amino acid-based, *e.g.*, sequence variants of OP-1, or antibody-derived sequences capable of functionally mimicking OP-1 binding to an OP-1-specific receptor. Examples of such antibodies may include anti-idiotypic antibodies. In a specific embodiment, the anti-idiotypic antibody mimics OP-1 both in receptor binding and in

- 8 -

ability to induce a transmembrane effect. Alternatively, the OP-1 analogs can be composed in part or in whole of other chemical structures, *e.g.*, the analogs can be comprised in part or in whole of nonproteinaceous molecules. In addition, the OP-1 analogs contemplated can be naturally-sourced or synthetically-produced.

5           As used herein, an OP-1 related protein contemplates a protein having a sequence sharing at least 60%, preferably greater than 65% or even 70% amino acid sequence homology or "similarity" with the C-terminal 96 amino acid sequence of OP-1 (OPS) as defined in Seq. ID Nos. 3-4, and which is able to substitute for OP-1 in ligand binding to ALK-1, (*e.g.*, able to compete successfully with OP-1 for binding to an ALK-1 receptor.) Proteins sharing at least 60%  
10 homology with OP-1 include BMP-2, BMP-4, and GDF-1. In another embodiment an OP-1-related protein is contemplated to include a protein having a sequence sharing at least 60% amino acid sequence homology with the C-terminal 102 amino acid sequence of OP-1, (OP-1 "seven cysteine domain"). In one preferred embodiment OP-1 related protein sequences are contemplated to include sequences sharing at least 60%, preferably greater than 65% or even  
15 70% identity with the C-terminal 102 amino acid sequence of OP-1. OP-1 related sequences contemplated by the invention include xenogenic homologs (*e.g.*, the *Drosophila* homolog 60A), and the related sequences referenced herein and in the literature as OP-2, OP-3, BMP-5, BMP-6 (and its xenogenic homolog Vgr-1.) OP-1 related sequences also include sequence variants encoded by a nucleic acid which hybridizes with a DNA sequence comprising the C-terminal 102  
20 amino acids of Seq. ID No. 3 under stringent hybridization conditions and which can substitute for OP-1 in an OP-1-receptor binding assay. In another embodiment, OP-1 sequence variants include proteins which can substitute for OP-1 in a ligand-receptor binding assay and which is recognized by an antibody having binding specificity for OP-1.

          As used herein, "amino acid sequence homology" is understood to mean amino acid  
25 sequence similarity, and homologous sequences sharing identical or similar amino acids, where similar amino acids are conserved amino acids as defined by Dayoff *et al.*, *Atlas of Protein Sequence and Structure*; vol. 5, Suppl. 3, pp. 345-362 (M.O. Dayoff, ed., Nat'l BioMed. Research Fdn., Washington D.C. 1978.) Thus, a candidate sequence sharing 60% amino acid homology with a reference sequence requires that, following alignment of the candidate sequence with the  
30 reference sequence, 60% of the amino acids in the candidate sequence are identical to the corresponding amino acid in the reference sequence, or constitute a conserved amino acid change

- 9 -

thereto. "Amino acid sequence identity" is understood to require identical amino acids between two aligned sequences. Thus, a candidate sequence sharing 60% amino acid identity with a reference sequence requires that, following alignment of the candidate sequence with the reference sequence, 60% of the amino acids in the candidate sequence are identical to the corresponding amino acid in the reference sequence.

As used herein, all OP-1-related protein homologies and identities use OP-1 as the reference sequence, with the C-terminal 102 amino acids described in Seq. ID Nos. 3-4 constituting the seven cysteine domain. Also as used herein, sequences are aligned for homology and identity calculations using the method of Needleman *et al.* ((1970) *J. Mol. Biol.* 48: 443-453), and identities are calculated by the Align program (DNASTar, Inc.). In all cases, internal gaps and amino acid insertions in the candidate sequence as aligned are ignored when making the homology/identity calculation.

In one aspect, the invention contemplates isolated ligand-receptor complexes comprising OP-1 or an OP-1 related protein or analog as the ligand in specific binding interaction with an ALK-1 receptor or receptor analog, as defined herein. In another aspect, the invention contemplates that the ligand-receptor complex also comprises part or all of an OP-1 binding Type II receptor. Type II receptors contemplated to be useful include Type II receptors defined in the literature as having binding specificity for activin or a bone morphogenic protein, including daf-4, ActRII and AtrII. In still another aspect, the ligand-receptor complex comprises both a Type I and a Type II receptor and OP-1, or an OP-1 related protein or analog as the ligand. In all complexes, the bound receptor may comprise just the extracellular, ligand binding domain, or may also include part or all of the transmembrane sequence, and/or the intracellular kinase domain. Similarly, the ligand may comprise just the receptor binding sequence, longer sequences, including the mature dimeric species or any soluble form of the protein or protein analog.

In another aspect, the invention contemplates binding partners having specific binding affinity for an epitope on the ligand-receptor complex. In a preferred embodiment, the binding partner can discriminate between the complex and the uncomplexed ligand or receptor. In another embodiment, the binding partner has little or no substantial binding affinity for the uncomplexed ligand or receptor. In another preferred embodiment, the binding partner is a binding protein, more preferably an antibody. These antibodies may be monoclonal or polyclonal,

may be intact molecules or fragments thereof (*e.g.*, Fab, Fab', (Fab')<sub>2</sub>), or may be biosynthetic derivatives, including, but not limited to, for example, monoclonal fragments, such as single chain F<sub>V</sub> fragments, referred to in the literature as sF<sub>V</sub>s, BABs and SCAs, and chimeric monoclonals, in which portions of the monoclonals are humanized (excluding those portions involved in antigen recognition (*e.g.*, complementarity determining regions, "CDRs".) See, for example, U.S. Pat. Nos. 5,091,513 and 5,132,405, the disclosures of which are incorporated herein by reference. Biosynthetic chimeras, fragments and other antibody derivatives may be synthesized using standard recombinant DNA methodology and/or automated chemical nucleic acid synthesis methodology well described in the art and as described below.

In still another aspect, the invention provides molecules useful in the design and/or identification of morphogenic protein analogs as described below, as well as kits and methods, *e.g.*, screening assays, for identifying these analogs. The molecules useful in these assays can include part or all of the ALK-1 receptor sequence of Seq. ID No. 1, including analogs and amino acid sequence variants thereof. Currently preferred for these assays are receptor sequences comprising at least the sequence which defines the extracellular, ligand binding domains. Similarly, the kits and screening assays can be used in the design and/or identification of ALK-1 receptor analogs. The OP-1 analogs and OP-1 receptor analogs identified then can be produced in reasonable quantities using standard recombinant expression or chemical synthesis technology.

In still another aspect, the receptor and/or OP-1-specific receptor analogs can be used in standard methodologies for affinity purifying and/or quantifying OP-1, OP-1-related proteins and analogs. For example, the receptor's ligand binding domain first may be immobilized on a surface of a well or a chromatographic column; ligand in a sample fluid then may be provided to the receptor under conditions to allow specific binding; non-specific binding molecules then removed, *e.g.*, by washing, and the bound ligand then selectively isolated and/or quantitated. Similarly, OP-1, OP-1-related proteins and OP-1 analogs can be used for affinity purifying and/or quantifying ALK-1 receptors and receptor analogs. In one embodiment, the method is useful in kits and assays for diagnostic purposes which detect the presence and/or concentration of OP-1 protein or related morphogen in a body fluid sample including, without limitation, serum, peritoneal fluid, spinal fluid, and breast exudate. The kits and assays can also be used for detecting and/or quantitating ALK-1 receptors in a sample.

- 11 -

In still another aspect the invention comprises ALK-1 receptors and receptor analogs useful in screening assays to identify organs, tissues and cell lines which express ALK-1 receptors. These cells then can be used in screening assays to identify ligands that modulate endogenous morphogen receptor expression levels, including the density of receptors expressed on a cell surface. Useful assay methodologies may be modeled on those described in PCT US92/07359, and as described below.

The invention thus relates to compositions and methods for the use of morphogen receptor sequences in diagnostic, therapeutic and experimental procedures. Active receptors useful in the compositions and methods of this invention can include truncated or full length forms, as well as forms having varying glycosylation patterns. Active receptors useful in the invention also include chimeric constructs as described below. Active OP-1-specific receptors/analogues can be expressed from intact or truncated genomic or cDNA, or from synthetic DNAs in prokaryotic or eukaryotic host cells, and purified, cleaved, refolded and oxidized as necessary to form active molecules. Useful host cells include prokaryotes, including *E. coli* and *B. subtilis*, and eukaryotic cells, including mammalian cells, such as fibroblast 3T3 cells, CHO, COS, melanoma or BSC cells, Hela and other human cells, the insect/baculovirus system, as well as yeast and other microbial host cell systems.

Thus, in view of this disclosure, skilled genetic engineers now can, for example, identify and produce ALK-1 cell surface receptors or analogs thereof; create and perform assays for screening candidate OP-1 or OP-1-related protein analogs and evaluate promising candidates in therapeutic regimes and preclinical studies; modulate the availability of endogenous morphogen for cell surface interactions; modulate endogenous ALK-1 cell surface receptor levels; elucidate the signal transduction pathway induced or otherwise mediated by ALK-1 receptor binding; and modulate tissue morphogenesis *in vivo*.

## Brief Description of the Drawings

Fig 1. is a schematic representation of OP-1-induced PAI-1 production in cells transfected with ALK-1, ALK-3 or ALK-5;

- 12 -

Fig 2A is a graphic representation of  $^3\text{H}$ -thymidine incorporation in response to OP-1 in R4-2 cells transfected with ALK-1 (closed circles), ALK-3 (closed triangles) and ALK-5 (closed squares) as a function of OP-1 concentration;

Fig 2B is a graphic representation of  $^3\text{H}$ -thymidine incorporation in response to OP-1 or TGF- $\beta$  in cells transfected with ALK-1, ALK-3 or ALK-5.

### Detailed Description

Disclosed herein is a Type I receptor, ALK-1, having binding specificity for true tissue morphogenic proteins, particularly OP-1 and OP-1-related proteins. The ALK-1 Type I receptors disclosed herein, can be used together with OP-1, OP-1-related proteins and OP-1 analogs for therapeutic, diagnostic and experimental uses as described herein below. Moreover, soluble forms of the receptor proteins, *e.g.*, forms consisting essentially of the extracellular domain or a fragment thereof sufficient to bind OP-1 with specificity, can be used as a soluble therapeutic morphogen antagonist, as described below.

Following this disclosure, related ALK-1 receptors are available, as are high and medium flux screening assays for identifying ligand analogs and ALK-1 receptor analogs. These analogs can be naturally occurring molecules, or they can be designed and biosynthetically created using a rational drug design and an established structure/function analysis. The analogs can be amino acid-based or can be composed in part or whole of non-proteinaceous synthetic organic molecules. Useful analogs also can include antibodies, preferably monoclonal antibodies (including fragments thereof, *e.g.*, Fab, Fab', and (Fab')<sub>2</sub>), or synthetic derivatives thereof, such as monoclonal single chain F<sub>v</sub> fragments known in the art as sF<sub>v</sub>s, BABs, and SCAs (see below), and bispecific antibodies or derivatives thereof. When these antibodies mimic the binding activity of OP-1 to a cell surface receptor without inducing the biological response OP-1 does upon binding, the antibody can compete for OP-1 binding and act as an antagonist. These antibodies or derivatives thereof can also mimic OP-1 both in receptor binding and signal transduction, in which case the antibody acts as an OP-1 agonist. The antibodies and derivatives can also be used for inducing the morphogenic cellular response by crosslinking receptors to morphogenic proteins, particularly OP-1 and OP-1-related proteins to form either homo- or hetero-complexes of the Type I and Type II receptors.

- 13 -

The ALK-1 receptor sequences described herein also can be used to create chimeric sequences, wherein, for example, part or all of either the extracellular domain or the intracellular domain is a non-ALK sequence or is created from two or more ALK sequences. These chimeric receptors can be synthesized using standard recombinant DNA methodology and/or automated chemical nucleic acid synthesis methodology well described in the art and as disclosed below. Chimerics can be used, for example, in OP-1 analog assays, wherein the OP-1-binding extracellular domain is coupled to a non-ALK intracellular domain that is well characterized and/or readily detectable as a second messenger response system, as described below. Chimerics can also be used, for example, in high flux OP-1 analogs screens and as part of purification protocols, wherein a soluble ligand binding domain of an OP-1-specific receptor is immobilized on to a support *e.g.*, by covalent or non-covalent interactions, with a chromatographic matrix or the well surface of a 96-well plate. When immobilized onto a chromatographic matrix surface, the receptor fragment can be used in a protocol to isolate OP-1 or OP-1 analogs. When immobilized on a well surface the receptor fragment is particularly useful in a screening assay to identify receptor-binding OP-1 analogs in a standard competition assay.

The true tissue morphogenic proteins contemplated to be useful in the methods and compositions of the invention include forms having varying glycosylation patterns and varying N-termini. The proteins can be naturally occurring or biosynthetically derived, and can be produced by expression of recombinant DNA in prokaryotic or eukaryotic host cells. The proteins are active as a single species (*e.g.*, as homodimers), or combined as a mixed species. Useful sequences and eukaryotic and prokaryotic expression systems are well described in the art. See, for example, US Patent Nos. 5,061,911 and 5,266,683 for useful expression systems.

Contemplated herein are OP-1 and OP-1-related sequences. Particularly useful OP-1 sequences are recited in US Pat Nos. 5,011,691; 5,018,753 and 5,266,683; in Ozkaynak *et al.* (1990) *EMBO J* 9: 2085-2093; and Sampath *et al.* (1993) *PNAS* 90: 6004-6008. OP-1 related sequences include xenogenic homologs, *e.g.*; 60A, from *Drosophila*, Wharton *et al.* (1991) *PNAS* 88: 9214-9218; and proteins sharing at least 60% amino acid sequence homology or "similarity" with OP-1 in the C-terminal seven cysteine domain. In a preferred embodiment, the related proteins share greater than 60% identity in the C-terminal seven cysteine domain, preferably at least 65% identity. Examples of OP-1 related sequences include BMP-5, BMP-6 (and its species homolog Vgr-1, Lyons *et al.* (1989) *PNAS* 86: 4554-4558), Celeste, *et al.* (1990) *PNAS* 87:

- 14 -

9843-9847 and PCT international application WO 93/00432; OP-2 (Ozkaynak *et al.* (1992) *J. Biol. Chem.* 267: 13198-13205) and OP-3 (PCT international application WO 94/06447). As will be appreciated by those having ordinary skill in the art, chimeric constructs readily can be created using standard molecular biology and mutagenesis techniques combining various portions of  
 5 different morphogenic protein sequences to create a novel sequence, and these forms of the protein also are contemplated herein.

A particularly preferred embodiment of the proteins contemplated by the invention includes proteins whose amino acid sequence in the cysteine-rich C-terminal domain has greater than 60% identity, and preferably greater than 65% identity with the amino acid sequence of OPS  
 10 (OP-1 sequence defining the C-terminal conserved six cysteines, *e.g.*, residues 335-431 of Seq. ID Nos. 3-4).

In another preferred aspect, the invention contemplates osteogenic proteins comprising species of polypeptide chains having the generic amino acid sequence herein referred to as "OPX" which accommodates the homologies between the various identified species of the osteogenic  
 15 OP-1 and OP-2 proteins, and which is described by the amino acid sequence presented below and in Seq. ID No. 5.

	Cys	Xaa	Xaa	His	Glu	Leu	Tyr	Val	Ser	Phe
	1				5					10
20	Xaa	Asp	Leu	Gly	Trp	Xaa	Asp	Trp	Xaa	Ile
					15					20
	Ala	Pro	Xaa	Gly	Tyr	Xaa	Ala	Tyr	Tyr	Cys
					25					30
25	Glu	Gly	Glu	Cys	Xaa	Phe	Pro	Leu	Xaa	Ser
					35					40
	Xaa	Met	Asn	Ala	Thr	Asn	His	Ala	Ile	Xaa
					45					50
	Gln	Xaa	Leu	Val	His	Xaa	Xaa	Xaa	Pro	Xaa
					55					60
30	Xaa	Val	Pro	Lys	Xaa	Cys	Cys	Ala	Pro	Thr
					65					70
	Xaa	Leu	Xaa	Ala	Xaa	Ser	Val	Leu	Tyr	Xaa
					75					80
	Asp	Xaa	Ser	Xaa	Asn	Val	Ile	Leu	Xaa	Lys
35					85					90
	Xaa	Arg	Asn	Met	Val	Val	Xaa	Ala	Cys	Gly
					95					100
	Cys	His,								



- 15 -

and wherein Xaa at res. 2 = (Lys or Arg); Xaa at res. 3 = (Lys or Arg); Xaa at res. 11 = (Arg or Gln); Xaa at res. 16 = (Gln or Leu); Xaa at res. 19 = (Ile or Val); Xaa at res. 23 = (Glu or Gln); Xaa at res. 26 = (Ala or Ser); Xaa at res. 35 = (Ala or Ser); Xaa at res. 39 = (Asn or Asp); Xaa at  
 5 res. 41 = (Tyr or Cys); Xaa at res. 50 = (Val or Leu); Xaa at res. 52 = (Ser or Thr); Xaa at res. 56 = (Phe or Leu); Xaa at res. 57 = (Ile or Met); Xaa at res. 58 = (Asn or Lys); Xaa at res. 60 = (Glu, Asp or Asn); Xaa at res. 61 = (Thr, Ala or Val); Xaa at res. 65 = (Pro or Ala); Xaa at res. 71 = (Gln or Lys); Xaa at res. 73 = (Asn or Ser); Xaa at res. 75 = (Ile or Thr); Xaa at res. 80 = (Phe or Tyr); Xaa at res. 82 = (Asp or Ser); Xaa at res. 84 = (Ser or Asn); Xaa at res. 89 = (Lys  
 10 or Arg); Xaa at res. 91 = (Tyr or His); and Xaa at res. 97 = (Arg or Lys).

In still another preferred aspect, the invention contemplates osteogenic proteins encoded by nucleic acids which hybridize to DNA or RNA sequences encoding the C-terminal seven cysteine domain of OP-1 or OP-2 under stringent hybridization conditions.

A brief description of the various terms of OP-1 useful in the invention is described below:

15 **OP-1** – Refers generically to the family of osteogenically active proteins produced by expression of part or all of the hOP-1 gene. Also referred to in related applications as "OPI" and "OP-1".

**OP-1-PP** – Amino acid sequence of human OP-1 protein (prepro form), Seq. ID Nos. 3-4, residues 1-431. Also referred to in related applications as "OP-1-PP" and "OPP".

20 **OP-1-18Ser** – Amino acid sequence of mature human OP-1 protein, Seq. ID Nos. 3-4, residues 293-431. N-terminal amino acid is serine. Originally identified as migrating at 18 kDa on SDS-PAGE in COS cells. Depending on protein glycosylation pattern in different host cells, also migrates at 23 kDa, 19 kDa and 17 kDa on SDS-PAGE. Also referred to in related applications as "OP-1-18."

25 **OP-1-16Ser; OP-1-16Ala; OP-1-16 Met; OP-1-16Leu; OP-1-16Val** – N-terminally truncated mature human OP-1 protein species defined, respectively, by residues 300-431; 316-431; 315-431; 313-431 and 318-431 of Seq. ID Nos. 3-4.

**OPS** – Amino acid sequence defining the C-terminal six cysteine domain, residues 335-431 of Seq. ID Nos. 3-4.

**OP7** – Amino acid sequence defining the C-terminal seven cysteine domain, residues 330-431 of Seq. ID Nos. 3-4.

**Soluble form OP-1** – mature dimeric OP-1 species having one or, preferably two copies of pro domain, *e.g.*, at least residues 158-292 of Seq. ID Nos. 3-4, preferably residues 48-292 or 30-292, non-covalently complexed with the dimer.

The cloning procedure for obtaining OP-1-binding ALK nucleic acid sequences, means for expressing receptor sequences, as well as other material aspects concerning the nature and utility of these sequences, including how to make and how to use the subject matter claimed, will be further understood from the following, which constitutes the best mode currently contemplated for practicing the invention.

#### **Example 1. IDENTIFICATION OF ALK-1**

The cloning and characterization of ALK-1, -2, -3, and -6 receptors are described in detail in ten Dijke *et al.* (1993) *Oncogene* 8: 2879-2887; and (1994) *Science* 264: 101-104. These molecules have similar domain structures: an N-terminal predicted hydrophobic signal sequence (von Heijne (1986) *Nucl. Acids Res.* 14: 4683-4690) is followed by a relatively small extracellular cysteine-rich ligand binding domain, a single hydrophobic transmembrane region (Kyte & Doolittle (1982) *J. Mol. Biol.* 157: 105-132) and a C-terminal intracellular portion, which consists almost entirely of a kinase domain.

The extracellular domains of the ALK-1 receptor is defined essentially by residues 22-118 (Seq. ID Nos. 1-2).

The positions of many of the cysteine residues in these receptors can be aligned, indicating that the extracellular domains likely adopt a similar structural configuration.

The intracellular domains of these receptors are characterized by a serine/threonine kinase, defined essentially by residues 204-494 (Seq. ID Nos. 1-2) for ALK-1. The catalytic domains of kinases can be divided into 12 subdomains with stretches of conserved amino-acid residues. The key motifs are found in serine/threonine kinase receptors indicating that they are functional kinases. The consensus sequence for the binding of ATP (Gly-X-Gly-X-X-Gly in subdomain I followed by a Lys residue further downstream in subdomain II) is found in all the ALK. Moreover, ALK-1, ALK-2, ALK-3 and ALK-6 have the sequence motifs or similar motifs

- 17 -

HRDLKSKN (Subdomain VIB) and GTKRYMAPE (Subdomain VIII), that are found in most of the serine/threonine kinase receptors and can be used to distinguish them from tyrosine kinase receptors. Two short inserts in the kinase domain (between subdomain VIA and VIB and between X and XI are unique to members of this serine/threonine kinase receptor family. In the intracellular domain, these regions, together with the juxtamembrane part and C-terminal tail, are the most divergent between family members.

## Example 2. RECEPTOR EXPRESSION

### A. General Considerations

Receptor DNA, or a synthetic form thereof, can be inserted, using conventional techniques well described in the art (see, for example, Maniatis (1989) *Molecular Cloning A Laboratory Manual*), into any of a variety of expression vectors and transfected into an appropriate host cell to produce recombinant protein polypeptide chains, including both full length and truncated forms thereof. Shortened sequences, for example, can be used for the production of soluble receptor fragments.

Useful host cells include *E. coli*, *Saccharomyces cerevisiae*, *Pichia pastoris*, the insect/baculovirus cell system, myeloma cells, and various other mammalian cells. The full length forms of the proteins of this invention preferably are expressed in mammalian cells, as disclosed herein. Soluble forms may be expressed from both mammalian or bacterial cell systems. The vector additionally may include various sequences to promote correct expression of the recombinant protein, including transcription promoter and termination sequences, enhancer sequences, preferred ribosome binding site sequences, preferred mRNA leader sequences, preferred protein processing sequences, preferred signal sequences for protein secretion, and the like. The DNA sequence encoding the gene of interest also may be manipulated to remove potentially inhibiting sequences or to minimize unwanted secondary structure formation. The recombinant morphogen receptor also may be expressed as a fusion protein. After translation, the protein may be purified from the cells themselves or recovered from the culture medium. The DNA also may include sequences which aid in expression and/or purification of the recombinant protein. One useful sequence for example, is a hexa-His (His<sub>6</sub>) sequence, which adds a histidine tail to allow affinity purification of the protein on a Cu<sup>2+</sup> IMAC column (see below.)

- 18 -

For example, the DNA encoding the extracellular domain may be inserted into a suitable expression vector for transformation into a prokaryote host such as *E. coli* or *B. subtilis*, to produce a soluble, morphogen binding fragment. The DNA may be expressed directly or may be expressed as part of a fusion protein having a readily cleavable fusion junction. An exemplary protocol for prokaryote expression using MR-1 DNA is provided below. Recombinant protein is expressed in inclusion bodies and may be purified therefrom using the technology disclosed in U.S. Patent No. 5,013,653, for example.

The DNA also may be expressed in a suitable mammalian host. Useful hosts include fibroblast 3T3 cells, (e.g., NIH 3T3, from CRL 1658) COS (simian kidney ATCC, CRL-1650) or CHO (Chinese hamster ovary) cells (e.g., CHO-DXB11, from Lawrence Chasin, (1980) *PNAS* 77: 4216-4222), mink-lung epithelial cells (MV1Lu), human foreskin fibroblast cells, human glioblastoma cells, and teratocarcinoma cells. Other useful eukaryotic cell systems include yeast cells, the insect/baculovirus system or myeloma cells.

To express an ALK-1 cell surface receptor, the DNA is subcloned into an insertion site of a suitable, commercially available vector along with suitable promoter/enhancer sequences and 3' termination sequences. Useful promoter/enhancer sequence combinations include the CMV promoter (human cytomegalovirus (MIE) promoter) present, for example, on pCDM8, as well as the mammary tumor virus promoter (MTV) boosted by the Rous sarcoma virus LTR enhancer sequence (e.g., from Clontech, Inc., Palo Alto). Expression also may be further enhanced using transactivating enhancer sequences. The plasmid also preferably contains an amplifiable marker, such as DHFR under suitable promoter control, e.g., SV40 early promoter (ATCC #37148). Transfection, cell culturing, gene amplification and protein expression conditions are standard conditions, well known in the art, such as are described, for example in Ausubel *et al.*, ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1989). Briefly, transfected cells are cultured in medium containing 5-10% dialyzed fetal calf serum (FCS), and stably transfected high expression cell lines obtained by amplification and subcloning and evaluated by standard Western and Northern blot analysis. Southern blots also can be used to assess the state of integrated receptor sequences and the extent of their copy number amplification.

The expressed protein then is purified using standard procedures. A currently preferred methodology uses an affinity column, such as a ligand affinity column or an antibody affinity

- 19 -

column, the bound material then washed, and receptor molecules selectively eluted in a gradient of increasing ionic strength, changes in pH or addition of mild denaturants. Alternatively, where a useful anchor sequence has been added to the DNA, such as a (His)<sub>6</sub> sequence, the column may be a standard affinity column such as Cu<sup>2+</sup> IMAC column. Here, for example, the cell culture media containing the recombinant protein is passed over a Cu<sup>2+</sup> IMAC column (for example, prepared with 25 mM imidazole). The bound protein then is washed with a compatible solution and eluted with EDTA. The anchor sequence can be removed by a standard chemical or enzymatic procedure.

Mammalian cell expression is preferred where morphogen receptor expression on a cell surface is desired. For example, cell surface expression may be desired to test morphogen or morphogen analog binding specificity for a cell surface receptor under *in vivo* conditions. Cell surface expression also may be most efficacious for medium flux cellular screen assays as described below.

### B.1 Exemplary Mammalian Cell Culture

The receptors described herein and tested in Example 8 described below, were expressed in (1) COS-1 cells; (2) mink lung epithelial cells (Mv1Lu); and (3) ATDC5, a chondrogenic cell line, (Atrami (1990) *Cell Diff. Dev.* 30: 109-116). They also can be tested in a variety of other cells including, without limitation, AG1518 human foreskin fibroblasts; MG-63 human osteosarcoma cells; PC12 rat pheochromocytoma cells (all obtainable from American Type Culture Collection, Rockville, MD); human glioblastoma cells (U-1240 MG, Bengt Westermarck, *et al.* (1988) *Cancer Research* 48: 3910-3918); Tera-2 teratocarcinoma cells (clone 13, Thompson *et al.* (1984) *J Cell Sci* 72: 37-64); MC3T3-E1 cells (Sudo *et al.* (1983) *J. Cell Biol.* 96: 191-198, and ROS 17/2.8 rat osteosarcoma cells (Majeska *et al.* (1985) *Endocrinology* 116: 170-179. Exemplary cultury conditions for the ROS cells include culture in Ham's F12 medium containing 14 mM HEPES buffer, 2.5 mM L-glutamine, 1.1 mM CaCl<sub>2</sub>, 5% fetal bovine serum and antibiotics; for MC3T3-E1 cells, culture in a-MEM with 10% fetal bovine serum and antibiotics, and for Tera-2 cells culture in 5% CO<sub>2</sub> atmosphere at 37°C in a-MEM containing 10% fetal bovine serum, 100 units/ml of penicillin and 50 mg/ml of streptomycin, using tissue culture dishes pretreated with 0.1% swine skin gelatin (Sigma) in phosphate-buffered saline. Unless otherwise specified, in the examples provided herein, cells were cultured in DMEM containing 10% fetal bovine serum and antibiotics.

- 20 -

### Example 3. ANTIBODY PRODUCTION

#### A. General Considerations

Antibodies capable of specifically binding the receptor molecules, ligand molecules, or the ligand-receptor complex itself, useful as analogs and useful in immunoassays and in the immunopurification of morphogen receptors described may be obtained as described below.

Where antibodies specific to the ALK-1 receptors are desired, but which do not interfere with ligand binding, the antigenic sequence preferably comprises the juxtamembrane sequence. Where antibodies capable of competing for ligand binding are desired, the ligand binding domain may be used as the antigen source. Where antibodies to the complex are desired, the complex itself preferably is used as the antigenic sequence and candidate antibodies then tested for cross reactivity with uncomplexed ligand and receptors versus the ligand-receptor complex. Finally, bispecific antibodies may be used to complex ligand to a cell surface receptor (Type I or Type II) and/or to target an agent or ligand to cells or tissue expressing a Type I or Type II morphogen-specific receptor. Preferred bispecific antibody derived molecules are single chain binding sites described in US Pat No. 5,091,513 and 5,132,405, the disclosures of which are incorporated hereinabove by reference.

Antibodies useful as OP-1 analogs may be obtained using the receptor ligand binding domain as the immunogen source and testing receptor-binding analogs for their ability to compete with OP-1 in a competition binding assay. Similarly, where antibodies useful as OP-1-specific receptor analogs are desired, OP-1 is the immunogen source and the antibody tested in a competition assay with receptor protein.

Polyclonal antibodies specific for a morphogen receptor of interest may be prepared generally as described below. Each rabbit is given a primary immunization (*e.g.*, 500 mg) of antigen in 0.1% SDS mixed with 500 ml Complete Freund's Adjuvant. The antigen is injected intradermally at multiple sites on the back and flanks of the animal. The rabbit is boosted after a month with 500 mg of antigen in the same manner using incomplete Freund's Adjuvant. Test bleeds are taken from the ear vein seven days later. Two additional boosts and test bleeds are performed at monthly intervals until antibody against the antigenic sequence is detected in the serum using a standard Western blot. Then, the rabbit is boosted monthly with 100 mg/ml of antigen and bled (15 ml per bleed) at days seven and ten after boosting.

Similarly, monoclonal antibodies specific for a given morphogen receptor molecule of interest may be prepared as described below: A mouse is given two injections of the antigenic sequence. The protein preferably is recombinantly produced. Where it is desired that the antibody recognize an epitope on the morphogen binding surface of a receptor an antigenic fragment derived from the extracellular domain preferably is provided. The first injection contains 100 mg of antigen in complete Freund's adjuvant and is given subcutaneously. The second injection contains 50 mg of antigen in incomplete adjuvant and is given intraperitoneally. The mouse then receives a total of 230 mg of antigen in four intraperitoneal injections at various times over an eight month period. One week prior to fusion, the mouse is boosted intraperitoneally with antigen (e.g., 100 mg) and may be additionally boosted with an antigen-specific peptide conjugated to bovine serum albumin with a suitable crosslinking agent. This boost can be repeated five days (IP), four days (IP), three days (IP) and one day (IV) prior to fusion. The mouse spleen cells then are fused to commercially available myeloma cells at a ratio of 1:1 using PEG 1500 (Boehringer Mannheim, Germany), and the fused cells plated and screened for ALK-specific antibodies using ALK-1, ALK-2, ALK-3 or ALK-6 as antigen. The cell fusion and monoclonal screening steps readily are performed according to standard procedures well described in standard texts widely available in the art. (See, for example, *Guide to Protein Purification* Murray P. Deutscher, ed., Academic Press, San Diego, 1990.

#### B. Exemplary ALK-Specific Antisera

Antibodies used in the assays described herein were obtained as reported in Franzen *et al.* (1993) *Cell* 75: 885-892, and ten Dijla *et al.* (1994) *Science* 264: 101-104. Briefly, rabbit antisera against ALK-1 were made against synthetic peptides corresponding to the divergent, intracellular juxtamembrane parts: residues 119-141. Peptides were synthesized with an Applied Biosystems 430 A Peptide Synthesizer using t-butoxycarbonyl chemistry, and purified by reverse phase HPLC. The synthetic peptides were coupled to keyhole limpet hemocyanin (Calbiochem-Behring) using glutaraldehyde, as described by Gullick *et al.* (1985) *EMBO J* 4: 2869-2877. The coupled peptides then were mixed with Freund's adjuvant and used to immunize rabbits using standard methodologies.

#### Example 4. OP-1-RECEPTOR BINDING ASSAYS

Ligand binding specificity is determined by evaluating the ability of a receptor molecule to bind a specific ligand, and the ability of that ligand to compete against itself and other molecules

- 22 -

which bind the receptor. Useful ligands will have a binding affinity for a soluble morphogen receptor extracellular domain such that dissociation constant ( $K_d$ ) is less than about  $10^{-6}$  M, preferably less than  $5 \times 10^{-7}$  M. Where stronger binding interaction is desired, preferred affinities are defined by a  $K_d$  of  $10^{-8}$ - $10^{-9}$  M. OP-1-related proteins are expected to be able to bind with specificity to multiple different receptor molecules, although likely with differing affinities.

Ligand binding specificity can be assayed as follows, essentially following standard protocols well described in the art and disclosed, for example, in Legerski *et al.* (1992) *Biochem. Biophys. Res. Comm.* 183: 672-679 and Frakar *et al.*, (1978) *Biochem. Biophys. Res. Comm.* 80: 849-857. In the ligand binding assays, a ligand having a known, quantifiable affinity for a morphogen receptor molecule of interest is labeled, typically by radioiodination ( $^{125}\text{I}$ ), *e.g.*, by chromogenic or fluorogenic labeling, or by metabolic labeling, *e.g.*,  $^{35}\text{S}$ , and aliquots of cells expressing the receptor on their surface are incubated with the labeled ligand, in the presence of various concentrations of unlabelled potential competitor ligand. In the assays described in Examples 8 and 9, below, this competitor typically is the candidate morphogen analog or an aliquot from a broth or extract anticipated to contain a candidate morphogen analog.

Alternatively, a crosslinking agent may be used to covalently link the ligand to the bound receptor, and the crosslinked complex then immunoprecipitated with an antibody specific to the ligand, receptor, or complex.

A standard, exemplary protocol for determining binding affinity is provided below. Briefly, cells expressing a receptor on their cell surface are plated into 35 mM dishes and incubated for 48 hours in DMEM (Dulbecco's modified Eagle medium) plus 10% fetal calf serum. Purified morphogen, here, *e.g.*, OP-1, or an OP-1-analog is iodinated with  $\text{Na}^{125}\text{I}$  by chloramine T oxidation, preferably having a specific activity of about 50-100 mCi/mg, essentially following the protocol of Frolik *et al.* (1984) *J. Biol. Chem.* 595: 10995-11000. Labeled morphogen then is purified using standard procedures, *e.g.*, chromatographically. Plated cells then are washed twice with physiologically buffered saline in the presence of 0.1% BSA, and incubated at 22°C in the presence of BSA, buffer and labeled morphogen (1 ng) and various concentrations (*e.g.*, 0-10mg/ml) of unlabelled competitor, *e.g.*, unlabelled morphogen or candidate ligand analogs. Following binding, cells are washed three times with cold buffer, solubilized in 0.5 ml of 0.5 N NaOH, removed from the dish, and radioactivity determined by gamma or scintillation counter.



- 23 -

Data then are expressed as percent inhibition, where 100% inhibition of specific binding is the difference between binding in the absence of competitor and binding in the presence of a 100-fold molar excess of unlabelled morphogen. Binding parameters preferably are determined using a computer program such as LIGAND (Munsun *et al.* (1980) *Anal. Biochem.* 107: 220-259.)

5       Where the receptor cell surface binding domain is to be provided as a soluble protein, the assay can be performed in solution, most readily as an immunoprecipitation assay. In currently preferred assays the morphogen molecule is labeled and incubated with unlabelled receptor and candidate morphogen analogs. Receptor-specific antibody then is provided to the solution to precipitate the receptor-morphogen complex and the amount of labeled morphogen in the  
10 precipitated complex determined using standard detection means.

      Where the receptor or ligand is to be used in an affinity isolation protocol, the molecule preferably is immobilized on a surface, preferably a matrix surface over which sample fluid will flow, allowing the ligand of interest to bind, at letting nonbinding components pass through as effluent. The complex then can be removed intact or the ligand selectively removed with a desired  
15 eluant.

#### 4.1 Screening Assay Considerations

      In an analog screening assay, the currently preferred protocol is a standard competition or radioimmunoassay (RIA). Here the OP-1 is labeled and the relative binding affinity of a candidate OP-1 analog ligand in a sample is measured by quantitating the ability of the candidate (unlabelled  
20 ligand analog) to inhibit binding of the labeled ligand (competitor morphogen) by the receptor. In performing the assay, fixed concentrations of receptor and labeled morphogen are incubated in the absence and presence of unknown samples containing candidate ligands. Sensitivity can be increased by preincubating the receptor with candidate ligand before adding the labeled morphogen. After the labeled competitor has been added, sufficient time is allowed for adequate  
25 competitor binding, and then free and bound labeled morphogen are separated, and one or the other is measured. Useful morphogen labels include radioactive labels, chromogenic or fluorogenic labels, and conjugated enzymes having high turnover numbers, such as horseradish peroxidase, alkaline phosphatase, or b-galactosidase, used in combination with chemiluminescent or fluorogenic substrates. In the same manner, OP-1-specific receptor analogs can be assayed for  
30 their affinity for OP-1 in competition assays with labeled OP-1 specific receptors.

## 4.2 Diagnostic Assay Considerations

The ability to detect morphogens in solution provides a valuable tool for diagnostic assays, allowing one to monitor the level of morphogen free in the body, *e.g.*, in serum, urine, spinal or peritoneal fluid, breast exudate, and other body fluids.

5           For example, OP-1 is an intimate participant in normal bone growth and resorption. Thus, soluble OP-1 is expected to be detected at higher concentrations in individuals experiencing high bone formation, such as children, and at substantially lower levels in individuals with abnormally low rates of bone formation, such as patients with osteoporosis, aplastic bone disease, or osteopenia. Monitoring the level of OP-1 in serum thus provides a means for evaluating the  
10          status of bone tissue and bone homeostasis in an individual, as well as a means for monitoring the efficacy of a treatment to regenerate damaged or lost bone tissue.

          For serum assays, the serum preferably first is partially purified to remove some of the excess, contaminating serum proteins, such as serum albumin. Preferably the serum is extracted by precipitation in ammonium sulfate (*e.g.*, 45%) such that the complex is precipitated. Further  
15          purification can be achieved using purification strategies that take advantage of the differential solubility of soluble morphogen complex or mature morphogens relative to that of the other proteins present in serum. Further purification also can be achieved by chromatographic techniques well known in the art. The sample fluid then can be assayed for OP-1 using the OP-1-specific receptor(s) and binding assays as described herein.

20          Morphogens useful in the binding/screening assays contemplated herein include the soluble forms of the protein, *e.g.*, the mature dimeric species complexed with one or two copies of the pro domain, the mature dimeric species alone, and truncated forms comprising essentially just the C-terminal active domain.

### Example 5. TRANSMEMBRANE SIGNAL INDUCTION ASSAYS/OP-1 MIMETICS

25          The kinase activity of the intracellular domains of the ALK-1 receptors can be tested in an autophosphorylation assay as described by Mathews *et al.* (PCT/US92/03825, published November 26, 1992). Briefly, the DNA fragment encoding at least the intracellular kinase domain of an OP-1-specific receptor is subcloned into pGEX-2T (Smith *et al.* (1988) *Gene* 67: 31-40) to create a fusion protein between the putative kinase domain and glutathione  
30          S-transferase (GST). The plasmid is introduced into *E. coli* and the expressed fusion protein

- 25 -

purified using glutathione affinity chromatography. About 100-200 ng of fusion protein or purified GST then are incubated with 25 mCi ( $^{32}\text{P}$ ) ATP in 50 mM Tris, 10 mM  $\text{MgCl}_2$  buffer for 30 minutes at  $37^\circ\text{C}$ . Products then are analyzed by gel electrophoresis and autoradiography. The fusion protein, but not GST alone, becomes phosphorylated, indicating that the kinase domain is functional. Phosphoamino acid analysis then can be performed to determine the predominant amino acid being phosphorylated. Similar assays can be performed using similar fusion constructs expressed in mammalian cells.

Various signaling transduction assays are provided in Example 9, below. An assay can also be developed for testing kinase activity transduction upon ligand binding using a ligand-induced kinase activity assay known in the art. Here, the ability of OP-1 analog to induce phosphorylation upon binding to the receptor is tested.

See, for example, various assays for measuring ligand-induced kinase activity described by Accili *et al.* (1991) *J. Biol. Chem.* 266: 434-439 and Nakamura *et al.* (1992) *J. Biol. Chem.* 267: 18924-18928. For example, ligand-induced kinase activity (*e.g.*, receptor autophosphorylation) can be measured *in vitro* by incubating purified receptor in the presence and absence of ligand (here, OP-1 or OP-1 analog, *e.g.*,  $10^{-7}$  M) under conditions sufficient to allow binding of the ligand to the receptor, followed by exposure to  $^{32}\text{P}$ -ATP (*e.g.*, 100 mCi in the presence of 10 mM Tris-HCl (pH 7.6), 10 mM  $\text{MgCl}_2$ , 10 mM  $\text{MnCl}_2$ , 1 mM dithiothreitol, 0.15 M NaCl, 0.1% Triton X-100 and 3% glycerol) and the amount of phosphorylation measured, *e.g.*, by SDS polyacrylamide gel electrophoresis and autoradiography following immunoprecipitation with antiphosphoserine, antiphosphothreonine or antiphosphotyrosine antibody (*e.g.*, commercially available or made using standard antibody methodologies.) While a low level of autophosphorylation may be detected in the absence of ligand, incubation with ligand is anticipated to significantly increase (*e.g.*, 5-20 fold increase) the amount of phosphorylation detected.

In another assay for detecting ligand-induced receptor autophosphorylation, involving intact cells, receptor DNA is transfected into a suitable host cell, *e.g.*, a fibroblast, which then is grown under standard conditions to create a confluent monolayer of cells expressing the receptor on their cell surface. On the day of the experiment, cells are incubated with or without ligand (*e.g.*, OP-1 or OP-1 analog, *e.g.*,  $10^{-7}$  M) at  $37^\circ\text{C}$ , and then quickly washed with a "stopping solution" containing ATP (*e.g.*, 0.1 M NaF, 4 mM EDTA, 10 mM ATP, 10 mM sodium

- 26 -

orthovanadate, 4 mM sodium pyrophosphate). The cells then are frozen in a dry ice/ethanol bath, solubilized and the receptors immunoprecipitated, *e.g.*, with an antireceptor antibody, as described herein. The immune complexes then are segregated, washed, separated by gel electrophoresis using standard procedures and transferred to a membrane for Western blot analysis using standard procedures. Phosphorylation of the receptor then can be visualized by immunodetection with a suitable antibody (*e.g.*, antiphosphoserine, antiphosphothreonine or antiphosphotyrosine), as described above. The bound antibody (*e.g.*, bound antiphosphoserine, antiphosphothreonine or antiphosphotyrosine) then can be detected with <sup>125</sup>I labeled protein A, followed by autoradiography. The amount of phosphorylated receptor detected is anticipated to be significantly greater (5-20 fold increase) in receptors incubated with ligand than receptors exposed to ATP in the absence of ligand.

Ligand-induced receptor phosphorylation of exogenous substrates similarly can be assayed essentially using the methodology described herein. Here, a suitable substrate (*e.g.*, a synthetic polypeptide containing serine, threonine or tyrosine amino acids) is provided to the receptor following ligand exposure and prior to incubation with ATP. The substrate subsequently can be segregated by immunoprecipitation with an antibody specific for the substrate, and phosphorylation detected as described above. As for autophosphorylation, the amount of phosphorylated substrate detected following ligand incubation is anticipated to be greater than that detected for substrates exposed to receptors in the absence of ligand.

## Example 6. CHIMERIC RECEPTOR MOLECULES

Chimeric receptor molecules, *e.g.*, comprising an ALK or ALK analog extracellular and transmembrane region and, for example, part or all of an intracellular domain from another, different receptor or an intracellular domain from a different cell surface molecule, may be constructed using standard recombinant DNA technology and/or an automated DNA synthesizer to construct the desired sequence. As will be appreciated by persons skilled in the art, useful junctions include sequences within the transmembrane region and/or sequences at the junction of either the intracellular or the extracellular domains. Also envisioned are chimeras where the extracellular domain or the intracellular domains themselves are chimeric sequences.

Chimeric sequences are envisioned to be particularly useful in screening assays to determine candidate binding ligands (*e.g.*, OP-1 analogs, see below), where the non-receptor intracellular domain provides a suitable second messenger response system that is easy to detect.

- 27 -

Potentially useful other second messenger response systems include those which, when activated, induce phosphoinositide hydrolysis, adenylate cyclase, guanylate cyclase or ion channels.

Chimeric receptor molecules have particular utility in gene therapy protocols. For example, a population of cells expressing a chimeric morphogen receptor molecule on their surface and competent for expressing a desired phenotype can be implanted in a mammal at a particular tissue locus. By careful choice of the ligand binding domain used on these receptors a physician can administer to the individual a morphogen agonist capable of: (1) binding to the chimeric receptor alone and (2) stimulating the proliferation and/or differentiation of the implanted cells without affecting endogenous cell populations.

10 **Example 7. CONSIDERATIONS FOR IDENTIFYING OTHER OP-1 SPECIFIC RECEPTORS IN NUCLEIC ACID LIBRARIES**

Identification of ALK-1 allows one to identify other ALK-1 related morphogen receptor sequences in different species as well as in different tissues. ALK-1 sequences themselves can be used as a probe or the sequence may be modified to account for other potential codon usage (*e.g.*, human codon bias.) Currently preferred probe sequences are those which encode the receptor's extracellular domain.

Probes based on the nucleic acid sequence of Seq. ID No. 1 can be synthesized on commercially available DNA synthesizers, *e.g.* Applied Biosystems model 381A, using standard techniques, *e.g.* Gait, *Oligonucleotide Synthesis: A Practical Approach*, (IRL Press, Washington D.C., 1984). It is preferable that the probes are at least 8-50 bases long, more preferably 18-30 bases long. Probes can be labeled in a variety of ways standard in the art, *e.g.* using radioactive, enzymatic or colormetric labels as described, for example, by Berent *et al.*, (May/June 1985) *Biotechniques*: 208-220; and Jablonski *et al.*, (1986) *Nucleic Acids Research* 14: 6115-6128.

Preferably, low stringency conditions are employed when screening a library for morphogen receptor sequences using a probe derived from OP-1-binding receptor. Preferred ALK-specific probes are those corresponding to bases encoding the extracellular domain ("ECD"), or encoding a unique (nonhomologous) sequence within the cytoplasmic domain. Useful probes may be designed from bases encoding the juxtamembrane region, for example. The probe may be further modified to use a preferred species codon bias. Alternatively, probes derived from the serine/threonine kinase domain can be used to identify new members of the

- 28 -

receptor kinase family which can be screened for OP-1 binding using the methods described in Example 8.

For example, for a probe of about 20-40 bases a typical prehybridization, hybridization, and wash protocol is as follows: (1) prehybridization: incubate nitrocellulose filters containing the  
5 denatured target DNA for 3-4 hours at 55°C in 5xDenhardt's solution, 6xSSC (20xSSC consists of 175g NaCl, 88.2 g sodium citrate in 800 ml H<sub>2</sub>O adjusted to pH. 7.0 with 10 N NaOH), 0.1% SDS, and 100 mg/ml denatured salmon sperm DNA, (2) hybridization: incubate filters in prehybridization solution plus probe at 42°C for 14-48 hours, (3) wash; three 15 minutes washes in 6xSSC and 0.1% SDS at room temperature, followed by a final 1-1.5 minute wash in 6xSSC  
10 and 0.1% SDS at 55°C. Other equivalent procedures, e.g. employing organic solvents such as formamide, are well known in the art.

Alternatively, morphogen receptor-specific DNA can be amplified using a standard PCR (polymerase chain reaction) methodology such as the one disclosed herein, to amplify approximately 500 base pair fragments. As for the hybridization screening probes described  
15 above, the primer sequences preferably are derived from conserved sequences in the serine/threonine kinase domain. Examples of useful PCR amplifications are disclosed in ten Dijke, *et al.* (1993) *Oncogene* 8: 2879-2887 and (1994) *Science* 264: 101-104, and which also describe the isolation protocols for ALK-1.

### 7.1 Tissue Distribution of Morphogen Receptors

20 Determining the tissue distribution of OP-1-specific receptors can be used to identify tissue and cell sources which express these receptors, to identify new, related OP-1-specific receptor molecules, as well as to identify target tissues for OP-1-receptor interactions under naturally occurring conditions. The OP-1 specific receptor molecules (or their mRNA transcripts) readily are identified in different tissues using standard methodologies and minor modifications  
25 thereof in tissues where expression may be low. For example, protein distribution can be determined using standard Western blot analysis or immunohistological detection techniques, and antibodies specific to the morphogen receptor molecules of interest. Similarly, the distribution of morphogen receptor transcripts can be determined using standard Northern hybridization protocols and transcript-specific probes or by *in situ* hybridization.

- 29 -

Any probe capable of hybridizing specifically to a transcript, and distinguishing the transcript of interest from other related transcripts can be used. Because the morphogen receptors described herein likely share high sequence homology in their intracellular domains, the tissue distribution of a specific morphogen receptor transcript may best be determined using a probe specific for the extracellular domain of the molecule. For example, a particularly useful ALK-specific probe sequence is one derived from a unique portion of the 5' coding sequence, the sequence corresponding to the juxtamembrane region, or the 5' or 3' noncoding sequences. The chosen fragment then is labeled using standard means well known and described in the art and herein.

Using these receptor-specific probes, which can be synthetically engineered or obtained from cloned sequences, receptor transcripts can be identified and localized in various tissues of various organisms, using standard methodologies well known to those having ordinary skill in the art. A detailed description of a suitable hybridization protocol is described in Ozkaynak, *et al.*, (1991) *Biochem. Biophys. Res. Comm.* 179: 116-123, and Ozkaynak, *et al.* (1992) *J Biol. Chem.* 267: 25220-25227. Briefly, total RNA is prepared from various tissues (*e.g.*, murine embryo and developing and adult liver, kidney, testis, heart, brain, thymus, stomach) by a standard methodologies such as by the method of Chomczynski *et al.* ((1987) *Anal. Biochem* 162: 156-159) and described below. Poly(A)<sup>+</sup> RNA is prepared by using oligo (dT)-cellulose chromatography (*e.g.*, Type 7, from Pharmacia LKB Biotechnology, Inc.). Poly(A)<sup>+</sup> RNA (generally 15 mg) from each tissue is fractionated on a 1% agarose/formaldehyde gel and transferred onto a Nytran membrane (Schleicher & Schuell). Following the transfer, the membrane is baked at 80°C and the RNA is cross-linked under UV light (generally 30 seconds at 1 mW/cm<sup>2</sup>). Prior to hybridization, the appropriate probe is denatured by heating. The hybridization is carried out in a lucite cylinder rotating in a roller bottle apparatus at approximately 1 rev/min for approximately 15 hours at 37°C using a hybridization mix of 40% formamide, 5xDenhardts, 5xSSPE, and 0.1% SDS. Following hybridization, the non-specific counts are washed off the filters in 0.1xSSPE, 0.1% SDS at 50°C.

**Example 8. DEMONSTRATION THAT ALK-1 IS AN OP-1-BINDING RECEPTOR****Experimental Procedures*****Cell Culture***

A chondrogenic cell line, ATDC5 (Atsumi *et al.*, 1990 *Cell Diff. Div.* 30: 109-116) was  
5 cultured in a 1: 1 mixture of Dulbecco's modified Eagle's medium (DMEM) and Ham's F12  
containing 5% fetal bovine serum, 10 µg/ml bovine insulin (Sigma Co., Cincinnati) and 100  
units/ml penicillin. Cells from a chemically-mutagenized Mv1 Lu cell line (R mutant, clone 4-2  
(R4-2)) (Laiho *et al.*, 1990 *J. Biol. Chem.* 285: 18518-18524) also were used. R4-2 cells were  
cultured in DMEM containing 10% FBS and 100 units/ml penicillin in 5% CO<sub>2</sub> atmosphere at  
10 37°C.

***Preparation of Polyclonal Antibodies***

Antisera against ALK-1 to -6 were made against synthetic peptides corresponding to the  
intracellular juxtamembrane parts of the receptors as described herein.

***Stable Transfection of cDNA***

15 Stable expression plasmids of human ALK-1, ALK-3 (also referred to in the art as  
BMPR-1A) and ALK-5 (also referred to in the art as TBR-1) were generated by subcloning the  
corresponding full length cDNA into pMEP4 (Wrana *et al.*, 1992, *Cell* 71: 1003-1014), a Zn<sup>2+</sup>-  
inducible mammalian expression vector, using convenient restriction enzyme sites in the polylinker  
region of the vector. To generate stable transfectants, R4-2 cells were transfected with 10 µg  
20 each of plasmids by a calcium phosphate precipitation method using a mammalian transfection kit  
(Stratagene, La Jolla) as described by the manufacturer. After 2 days, selection was initiated by  
adding 120 U/ml hygromycin B (Wako Chemicals) to the culture medium. Independent colonies  
were cloned, and after screening by immunoprecipitation following the metabolic labeling of the  
cells, positive clones were chosen and further analyzed. More than two positive clones for each  
25 of the transfectants were subjected to the following experiments, in which no significant difference  
between independent clones was observed in the magnitude of receptor expression and response  
to the ligands.

***Recombinant Proteins and Radioiodination***

Recombinant human TGF-β1 and recombinant human activin A were obtained from H.  
30 Ohashi (Kirin Brewery Company Ltd.) and Y. Eto (Ajinomoto Company, Inc.), respectively.  
Recombinant human OP-1 was obtained as described (Sampath *et al.*, 1992). Iodination of OP-1



was performed according to the chloramine T method as described (ten Dijke *et al.*, (1994) *J. Biol. Chem.* 269: 16985-16988).

### ***Metabolic Labeling and Immunoprecipitation***

Metabolic labeling of stably transfected R4-2 cells was performed for 5 h in methionine- and cysteine-free DMEM with 100  $\mu\text{Ci}/\text{ml}$  of [ $^{35}\text{S}$ ]methionine and [ $^{35}\text{S}$ ]cysteine (Pro-mix cell labeling mix; Amersham) in the presence or absence of 100  $\mu\text{M}$   $\text{ZnCl}_2$ . After labeling, the cells were washed in phosphate-buffered saline and lysed in 1 ml of 20 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM EDTA, 1% Triton X-100, 1% sodium deoxycholate, 1.5% Trasylol (Bayer) and 1 mM phenylmethylsulfonyl fluoride (Sigma). The cell lysates were centrifuged, and supernatants were subjected to immunoprecipitation as previously described (Ichijo, *et al.*, (1993) *J. Biol. Chem.* 268: 14505-14513). Briefly, the metabolically labeled lysates were incubated with the antisera against each of ALKs for 45 min at 4°C. Immune complexes were bound to protein A-Sepharose (Kabi-Pharmacia) for 30 min at 4°C, washed four times with a buffer containing 20 mM Tris-HCl, pH 7.5, 500 mM NaCl, 1% Triton X-100, 1% deoxycholate, 0.2% sodium dodecyl sulfate (SDS), followed by one wash in distilled water. The immune complexes were eluted by boiling for 5 min in SDS-sample buffer (100 mM Tris-HCl, pH 8.8, 0.01% bromophenol blue, 36% glycerol, 4% SDS) containing 10 mM dithiothreitol and analyzed by 8.5% SDS-gel electrophoresis. The gel was fixed, dried and subjected to the analysis using a Fuji BAS 2000 Bio-Imaging Analyzer (Fuji Photo Film).

### ***Binding, Affinity Cross-Linking and Immunoprecipitation***

For receptor affinity labeling, ATDC5 cells were incubated on ice for 3 hrs with 300 pM of  $^{125}\text{I}$ -OP-1 in the presence or absence of 15 nM unlabeled OP-1 in the binding buffer (phosphate-buffered saline containing 0.9 mM  $\text{CaCl}_2$ , 0.49 mM  $\text{MgCl}_2$  and 1 mg/ml bovine serum albumin). For the ALK-transfected R4-2 cells, binding was allowed after overnight culture of the cells in the presence and absence of 100  $\mu\text{M}$   $\text{ZnCl}_2$ . After binding, cells were washed 3 times with the binding buffer and once with phosphate-buffered saline, and crosslinking was performed in the phosphate-buffered saline containing 0.28 mM of disuccinimidyl suberate (DSS; Pierce Chemical Co.) and/or 0.5 mM of Bis (sulfosuccinimidyl) suberate (BS3; Pierce) for 15 min on ice. Cells were washed once with the buffer containing 10 mM Tris-HCl, pH 7.4, 1 mM EDTA and 10% glycerol. Cell lysates were collected after incubation of the cells with 10 mM Tris-HCl pH 7.8, 1% NP-40, 0.15 M NaCl, 1 mM EDTA and 1.5% Trasylol for 20 min, and the supernatants were

- 32 -

analyzed by SDS-gel electrophoresis under reducing conditions or further subjected to immunoprecipitation. Immunoprecipitation of the labeled complex was done as described above. The immune complexes were eluted by boiling for 5 min in the SDS-sample buffer with 10 mM dithiothreitol, and analyzed using 7 or 8.5% polyacrylamide, followed by analysis using Bio-  
5 Imaging Analyzer.

#### ***PAI-1 Assay***

PAI-1 assay was performed as described previously (Ohtsuki and Massagué, (1992) *Mol. Cell. Biol.* 12: 261-265) with minor modifications. In brief, cells were seeded in 6-well cell culture plates and incubated with 100  $\mu$ M ZnCl<sub>2</sub> overnight. The cells were exposed to OP-1, TGF- $\beta$ 1 or activin A in serum-free DMEM without methionine and cysteine for 2 hrs. Thereafter,  
10 cultures were labeled with 30  $\mu$ Ci/ml of [<sup>35</sup>S]methionine and [<sup>35</sup>S]cysteine mixture for 2 hrs. The cells were removed by washing on ice once in phosphate-buffered saline, four times in 10 mM Tris-HCl, pH 8.0, 0.5% sodium deoxycholate and 1 mM phenylmethylsulphonyl fluoride, two times in 2 mM Tris-HCl, pH 8.0, and once in phosphate-buffered saline. Extracellular-matrix  
15 proteins were scraped off and extracted into SDS-sample buffer containing 10 mM dithiothreitol and analyzed by 10% SDS polyacrylamide gel electrophoresis, followed by analysis using Bio-imaging Analyzer. PAI-1 was identified as a 44 to 46 kDa doublet.

#### ***<sup>3</sup>H-Thymidine incorporation Assay***

Cells were seeded in 24-well cell culture plates at a density of 104 cells per well in DMEM  
20 with 10% FBS, and incubated overnight. The medium was changed to DMEM containing 0.2% FBS and 100  $\mu$ M ZnCl<sub>2</sub>, and the cells were incubated for 5 h. The medium was then changed to fresh DMEM containing 0.2% FBS, 100  $\mu$ M ZnCl<sub>2</sub> and various concentrations of OP-1 or TGF- $\beta$ 1. After 16 h of incubation, 0.25  $\mu$ Ci of <sup>3</sup>H-thymidine (Amersham) was added and the cells were incubated for an additional 2 h. Thereafter, the cells were fixed in 10% trichloroacetic  
25 acid for more than 15 min on ice, and solubilized with 1 M NaOH. The cell extracts were neutralized with 1 M HCl and <sup>3</sup>H radioactivity was determined in a liquid scintillation counter.

#### ***Identification of OP-1 Receptors in An OP-1 Responsive Cell Line.***

To identify the signaling receptor(s) for OP-1, we first attempted to find cells were identified which respond to OP-1. The mouse chondrogenic cell line ATDC5 was found to  
30 respond well to OP-1 and its proliferation and by alkaline phosphates activity were strongly stimulated. When the ATDC5 cells were analyzed for the binding of <sup>125</sup>I-OP-1 followed by

- 33 -

affinity cross-linking and SDS polyacrylamide gel electrophoresis under reducing conditions,  $^{125}\text{I}$ -OP-1 formed three major crosslinked complexes of 50-60 kDa, 70-85 kDa and 95-110 kDa which were effectively competed for binding in the presence of excess unlabeled OP-1. The size of the 70-85 and 95-110 complexes were similar to those of the type I and type II receptor complexes for TGF- $\beta$  and activin (ten Dijke *et al.*, (1994) *Science* 264: 101-104). The 50-60 kDa component may be a degradation product of a larger complex. In order to exclude the background and to investigate whether certain ALKs may serve as type I receptors for OP-1 in this particular cell line, the cross-linked complexes were further analyzed by immunoprecipitating these complexes using antisera against ALK-1 to ALK-6. Cross-linked complexes of 70-90 kDa could be precipitated by antisera to ALK-1 and ALK-3/BMPR-IA but not by others (*e.g.*, ALK-2, -4, -5, -6), indicating that ALK-1 as well as ALK-3/BMPR-IA form part of the isolated complexes and are competent to bind OP-1. ALK-1 may act as an endogenous Type I receptor for OP-1 or an OP-1-related protein.

***Expression and OP-1 Binding of ALKs in Stably Transfected R 4-2 Cells.***

Cells of a highly transfectable mink lung epithelial mutant cell line, R4-2, which lack endogenous type I receptor for TGF- $\beta$ . Were stably transfected with the ALK-1, ALK-3/BMPR-IA and ALK-5/TBR-I cDNAs under the control of Zn $^{2+}$ -inducible metallothionein promoter in an expression vector pMEP4. The R4-2 transfectants were metabolically labeled with [ $^{35}\text{S}$ ]methionine and [ $^{35}\text{S}$ ]cysteine in the presence or absence of 100  $\mu\text{M}$  ZnCl $_2$ , and the labeled proteins were immunoprecipitated by the antisera against each of ALKs. Labeled components of 50-65 kDa which fit with the sizes predicted from each of ALK cDNA sequences were specifically immunoprecipitated only in the presence of ZnCl $_2$ . A broad migration pattern of the components on the gel likely represents heterogenous glycosylation and/or phosphorylation.

In order to investigate whether ALK-1 and ALK-3/BMPR-IA can bind OP-1 when they were expressed in R4-2 cells, binding of  $^{125}\text{I}$ -OP-1 was analyzed by affinity cross-linking followed by immunoprecipitation to eliminate the background. When ALK-1 or ALK-3/BMPR-IA was induced by ZnCl $_2$ ,  $^{125}\text{I}$ -OP-1 bound to ALK-3/BMPR-IA, and to a lesser extent ALK-1, and the cross-linked complexes of 70-95 kDa were precipitated by the respective antisera. Intensity of the band for ALK-1 was weak, however, no cross-linked complex was detected in cells without Zn $^{2+}$  induction, or in ALK-5/TBR-1 expressing cells. When TGF- $\beta$ 1 was used as a radiolabeled ligand, only ALK-5/TBR-1 but not ALK-1 or ALK-3/BMPR-IA formed cross-linked complex

- 34 -

with  $^{125}\text{I}$ -TGF- $\beta$ 1. These results indicate that OP-1 and/or OP-1 related proteins can bind ALK-1 and ALK-3 when they are expressed in R4-2 cells. The results are presented in Table I below. Binding was specific as determined by standard competition assays. The values represented by "+/-", "+", "++", "+++", and "-" are all qualitative descriptors of the relative amount of radioactivity measured when the crosslinked molecules were gel electrophoresed and subjected to autoradiography. More radioactivity measured indicates a stronger binding interaction detected. In the Table the strength of binding interaction is as follows: +++ > ++ > + > +/- > -.

TABLE I

	$^{125}\text{I}$ -OP-1	$^{125}\text{I}$ -TGF- $\beta$
ALK-1	+	-
ALK-3	++	++
ALK-5	-	-

#### OP-1 Signals through ALK-1 and ALK-3/BMPR-IA

To investigate whether ALK-1 and ALK-3/BMPR-IA transduce any signals upon binding of OP-1, their abilities were determined to mediate signals for extracellular matrix production and cell growth production of PAI-1 was tested using the stable transfectants (Fig. 1). Production of PAI-1 in response to the stimulation by OP-1 in ALK-5/TBR-1 transfected cells was not detected. However, in ALK-1 and ALK-3/BMPR-IA-transfected cells, TGF- $\beta$ 1 and activin had no detectable effects on PAI-1 production, whereas OP-1 dramatically induced the PAI-1 protein in those cells in a  $\text{ZnCl}_2$  dependent manner. These results indicate that ALK-1 and ALK-3/BMPR-IA can signal an extracellular matrix production response upon binding of OP-1.

$^3\text{H}$ -thymidine incorporation assays were performed to investigate the ability of the three type I receptors ALK-1, ALK-3, ALK-5 to confer growth responsiveness to OP-1. When ALKs were induced in the presence of  $\text{ZnCl}_2$ , R4-2 cells expressing ALK-1 were found to respond to OP-1, and their growth was inhibited in a dose dependent manner (Fig. 2A). In contrast, OP-1 had no detectable effect on the cells expressing ALK-3/BMPR-IA or ALK-5/TBR-1 (Fig. 2A). In addition, the growth inhibitory effect of OP-1 on the ALK-1 transfected cells were dependent on the expression of ALK-1 since no inhibition of  $^3\text{H}$ -thymidine incorporation was observed in the

- 35 -

absence of  $\text{ZnCl}_2$ . (Fig. 2B). Under these conditions TGF- $\beta$  only inhibited cell growth in ALK-5 transfected cells.

The experimental results indicate that OP-1 binds ALK-1. The relatively weaker labeling intensity of ALK-1 by  $^{125}\text{I}$ -OP-1 as compared with ALK-3 in ATDC5 cells and transfected R4-2  
5 cells could be ascribed to reduced cross-linking efficiency with OP-1 since ALK-1 has fewer lysine residues in its extracellular domain (2 or 3, in human and mouse, respectively) than ALK-3 (7 lysines). Alternatively, the presence of a Type II receptor could be required for tight complex formation; and/or an OP-1 related protein could bind with higher affinity.

OP-1 has been shown to suppress cell proliferation and stimulate the expression of matrix  
10 proteins in rat osteosarcoma cells (Maliakal *et al.*, 1994, *Growth Factors* 11: 227-234). The data presented herein demonstrates that ALK-1 is competent to mediate growth-related signals in response to OP-1.

#### Example 9. OP-1, OP-1-SPECIFIC RECEPTOR ANALOG SCREENING ASSAYS

The present invention is useful to determine whether a ligand, such as a known or putative  
15 drug, is capable of binding to and/or activating an OP-1-specific cell surface receptor as described herein. Ligands capable of specific binding interaction with a given OP-1-specific receptor (*e.g.*, ALK-2, ALK-3, ALK-6) are referred to herein as OP-1 analogs and can be used for therapeutic and diagnostic applications. Some analogs will have the ability to stimulate morphogenic activity in the cell, mimicking both the receptor binding and signal transducing activity of OP-1. These  
20 are referred to OP-1 agonists or mimetics. Others will have strong binding affinity but will not stimulate morphogenesis, these are OP-1 antagonists. The analogs can be amino acid-based, or they can be composed of non-proteinaceous chemical structures.

The methods and kits described below similarly can be used to identify OP-1-specific receptor analogs, capable of mimicking the binding affinity of ALK-2, ALK-3 or ALK-6 for  
25 OP-1. The analogs can be provided to a mammal to interact with serum-soluble OP-1, effectively sequestering the protein and modulating its availability for cell surface interaction.

Transfection of an isolated clone encoding a morphogen receptor into the cell systems described above provides an assay system for the ability of ligands to bind to and/or to activate the receptor encoded by the isolated DNA molecule. Transfection systems, such as those  
30 described above, are useful as living cell cultures for competitive binding assays between known

- 36 -

or candidate drugs and ligands which bind to the receptor and compete with the binding of known morphogens, which are labeled by radioactive, enzymatic, spectroscopic or other reagents.

Membrane preparations containing the receptor and isolated from transfected cells are also useful in these competitive binding assays. Alternatively, and currently preferred, purified receptor

5 molecules or their ligand binding extracellular domains can be plated onto a microtiter well surface, in a modification of a sandwich assay, *e.g.*, as a competition assay, such as an RIA, described above. Finally, as described above, solution assays, and using only the receptor extracellular domain, also may be used to advantage in these assays. Functional assays of second messenger systems or their sequelae in transfection systems act as assays for binding affinity and  
10 efficacy in the activation of receptor function or efficacy in the antagonism of receptor function. Such a transfection system constitutes a "drug discovery system", useful for the identification of natural or synthetic compounds with potential for drug development that can be further modified or used directly as therapeutic compounds to activate or inhibit the natural functions of the receptor encoded by the isolated DNA molecule.

15 Once such candidate drugs (*e.g.*, OP-1 or receptor-binding analogs thereof) are identified, they can be produced in reasonable, useful quantities using standard methodologies known in the art. Amino acid-based molecules can be encoded by synthetic nucleic acid molecules, and expressed in a recombinant expression system as described herein above or in the art. Alternatively, such molecules can be chemically synthesized, *e.g.*, by means of an automated  
20 peptide synthesizer, for example. Non-amino acid-based molecules can be produced by standard organic chemical synthesis procedures. Where the candidate molecule is of undetermined structure, or composition, its composition readily can be determined by, for example, mass spectroscopy. Two approaches to identifying analogs typically are practiced in the art: high flux screens and rational design of ligand mimetics. High flux screens typically screen naturally  
25 sourced materials or chemical banks for their ability to bind a protein of interest, here, *e.g.*, the receptor. Typically, compounds are obtained from a range of sources, *e.g.*, chemical banks, microbial broths, plant and animal extracts, and the like. In a high flux screen typically, purified receptor, preferably the soluble, ligand binding extracellular domain, is plated onto a microtiter well surface and a standard volume of a sample solution to be tested then is added. Also added is  
30 a standard volume having a known quantity of a purified ligand known to bind the receptor with specificity. Preferably the ligand is labeled with a substance that is readily detectable by

- 37 -

automated means (e.g., radiolabel, chromophoric, fluorometric, enzymatic or spectroscopic label). The wells then are washed and the amount of label remaining after washing or the amount of label remaining associated with the receptor then is detected. Positive scores are identified by the ability of the test substance to prevent interaction of the labeled ligand with the receptor. The screening assays can be performed without undue experimentation, using standard molecular and cell biology tools in common use in the art. For example, screening assays can be performed in standard 96-well plates. Fifteen such plates reasonably can be set up at a time to perform multiple screening assays in parallel. Thus, with only 10-11 reiterations of the screening assay 15,625 (56) compounds can be screened for their binding affinity. Even allowing for a maximum incubation time of 2 hours, all 15,625 compounds reasonably can be assayed in a matter of days.

High flux screens exploit both the high degree of specificity of the labeled ligand for its receptor, as well as high throughput capacity of computer driven robotics and computer handling of data. Candidate analogs identified in this manner, then can be analyzed structurally and this information used to design and to synthesize analogs having enhanced potency, increased duration of action, increased selectivity and reduced side effects. Candidates also can be used in a rational design program as described below. Finally, candidate analogs also can be tested to determine morphogenic effect, if any, as described below.

The second approach to the identification of analogs uses a rational design approach to create molecules capable of mimicking the binding effect of OP-1 with an OP-1-specific receptor. Here the relevant structure for receptor binding is analyzed to identify critical sequences and structures necessary for binding activity and this information can be used to design and synthesize minimal size morphogen analogs. As for candidate compounds in the high flux assay, design candidates can be tested for receptor binding activity as described above. As described above, a candidate sequence can be further modified by, for example standard biological or chemical mutagenesis techniques to create a candidate derivative having, for example, enhanced binding affinity or another preferred characteristic.

Antibodies capable of interacting specifically with the receptor and competing with OP-1 binding can also be used as an analog. Antibodies can be generated as described above.

OP-1 analogs may be evaluated for their ability to mimic OP-1 or to inhibit OP-1 binding (e.g., agonists or antagonists) by monitoring the effect of the analogs on cells bearing an OP-1-

- 38 -

specific receptor (*e.g.*, ALK-1). OP-1 agonists are anticipated to have utility in any application where tissue morphogenesis is desired, such as in the regeneration of damaged tissue resulting from mechanical or chemical trauma, degenerative diseases, tissue destruction resulting from chronic inflammation, cirrhosis, inflammatory diseases, cancer and the like, and in the  
5 regeneration of tissues, organs and limbs. OP-1 antagonists are envisioned to have utility in applications where tissue morphogenesis is to be limited as, for example, in the treatment of malignant transformations including, but not limited to, osteosarcomas and Paget's disease.

Several exemplary systems for assaying the ability of a candidate analog transduce an OP-1-specific signal across the cellular membrane are described below.

#### 10 9.1 Induction of Osteoblast Differentiation Markers

For example, OP-1 is known to preferentially induce differentiation of progenitor cells, including embryonic mesenchymal cells and primary osteoblasts (see, for example, PCT US92/07432). As one example, OP-1 analogs can be tested for their ability to induce  
15 differentiation of primary osteoblasts, by measuring the ability of these analogs to induce production of alkaline phosphatase, PTH-mediated cAMP and osteocalcin, all of which are induced when primary osteoblasts are exposed to OP-1, 60A or dpp.

Briefly, the assays may be performed as follows. In this and all examples involving osteoblast cultures, rat osteoblast-enriched primary cultures preferably are used. Although these cultures are heterogeneous in that the individual cells are at different stages of differentiation,  
20 these cultures are believed to more accurately reflect the metabolism and function of osteoblasts *in vivo* than osteoblast cultures obtained from established cell lines. Unless otherwise indicated, all chemicals referenced are standard, commercially available reagents, readily available from a number of sources, including Sigma Chemical, Co., St. Louis; Calbiochem, Corp., San Diego and Aldrich Chemical Co., Milwaukee.

25 Rat osteoblast-enriched primary cultures are prepared by sequential collagenase digestion of newborn suture-free rat calvaria (*e.g.*, from 1-2 day-old animals, Long-Evans strain, Charles River Laboratories, Wilmington, MA), following standard procedures, such as are described, for example, in Wong *et al.*, (1975) *PNAS* 72: 3167-3171. Rat osteoblast single cell suspensions then are plated onto a multi-well plate (*e.g.*, a 24 well plate) at a concentration of 50,000 osteoblasts  
30 per well in alpha MEM (modified Eagle's medium, Gibco, Inc., Long Island) containing 10% FBS



- 39 -

(fetal bovine serum), L-glutamine and penicillin/streptomycin. The cells are incubated for 24 hours at 37°C, at which time the growth medium is replaced with alpha MEM containing 1% FBS and the cells incubated for an additional 24 hours so that the cells are in serum-deprived growth medium at the time of the experiment.

#### 5        ***Alkaline Phosphatase Induction of Osteoblasts***

The cultured cells in serum-free medium are incubated with OP-1, OP-1 analog or a negative control, using a range of concentrations. For example, 0.1, 1.0, 10.0, 40.0 or 80.0 ng OP-1/ml medium typically are used. 72 hours after the incubation period the cell layer is extracted with 0.5 ml of 1% Triton X-100. The resultant cell extract then, is centrifuged, and  
10    100 ml of the extract is added to 90 ml of paranitrosophenylphosphate (PNPP)/glycerine mixture and incubated for 30 minutes in a 37°C water bath and the reaction stopped with 100 ml NaOH. The samples then are run through a plate reader (e.g., Dynatech MR700 plate reader, and absorbance measured at 400 nm, using p-nitrophenol as a standard) to determine the presence and amount of alkaline phosphate activity. Protein concentrations are determined by the Biorad  
15    method. Alkaline phosphatase activity is calculated in units/mg protein, where 1 unit=1 nmol p-nitrophenol liberated/30 minutes at 37°C. OP-1 induces a 5-fold increase in the specific activity of alkaline phosphate by this method. Agonists are expected to have similar induction effects. Antagonists should inhibit or otherwise interfere with OP-1 binding, and diminished alkaline phosphatase induction should result when the assay is performed with an antagonist in the presence  
20    of a limiting amount of OP-1.

#### ***Induction of PTH-Mediated cAMP.***

The effect of a morphogen analog on parathyroid hormone-mediated cAMP production in rat osteoblasts *in vitro* may be demonstrated as follows.

Rat osteoblasts are prepared and cultured in a multiwell plate as described above. The  
25    cultured cells then are divided into three groups: (1) wells which receive, for example, 1.0, 10.0 and 40.0 ng OP-1/ml medium); (2) wells which receive the candidate analog at various concentration ranges; and (3) a control group which receives no additional factors. The plate is then incubated for another 72 hours. At the end of the 72 hours the cells are treated with medium containing 0.5% bovine serum albumin (BSA) and 1 mM 3-isobutyl-1-methylxanthine for  
30    20 minutes followed by the addition into half of the wells of human recombinant parathyroid

- 40 -

hormone (hPTH, Sigma, St. Louis) at a concentration of 200 ng/ml for 10 minutes. The cell layer then is extracted from each well with 0.5 ml of 1% Triton X-100. The cAMP levels then are determined using a radioimmunoassay kit (e.g., Amersham, Arlington Heights, Illinois). OP-1 doubles cAMP production in the presence of PTH. Agonists are expected to have similar induction effects. Antagonists are expected to inhibit or otherwise interfere with OP-1 binding, and diminished cAMP production should result when the assay is performed with an antagonist in the presence of limiting the amount of OP-1.

### *Induction of Osteocalcin Production*

Osteocalcin is a bone-specific protein synthesized by osteoblasts which plays an integral role in the rate of bone mineralization *in vivo*. Circulating levels of osteocalcin in serum are used as a marker for osteoblast activity and bone formation *in vivo*. Induction of osteocalcin synthesis in osteoblast-enriched cultures can be used to demonstrate morphogenic efficacy *in vitro*.

Rat osteoblasts are prepared and cultured in a multi-well plate as above. In this experiment the medium is supplemented with 10% FBS, and on day 2, cells are fed with fresh medium supplemented with fresh 10 mM  $\beta$ -glycerophosphate (Sigma, Inc.). Beginning on day 5 and twice weekly thereafter, cells are fed with a complete mineralization medium containing all of the above components plus fresh L(+)-ascorbate, at a final concentration of 50 mg/ml medium. OP-1 or OP-1 analog then is added to the wells directly, e.g., in 50% acetonitrile (or 50% ethanol) containing 0.1% trifluoroacetic acid (TFA), at no more than 5 ml OP-1/ml medium. Control wells receive solvent vehicle only. The cells then are re-fed and the conditioned medium sample diluted 1:1 in standard radioimmunoassay buffer containing standard protease inhibitors and stored at -20°C until assayed for osteocalcin. Osteocalcin synthesis is measured by standard radioimmunoassay using a commercially available osteocalcin-specific antibody and can be confirmed by Northern blot analysis to calculate the amount of osteocalcin mRNA produced in the presence and absence of OP-1 or OP-1 analog. OP-1 induces a dose-dependent increase in osteocalcin production (5-fold increase using 25 ng of OP-1 protein/ml), and a 20-fold increase in osteocalcin mRNA. Agonists are expected to have similar induction effects; antagonists are expected to inhibit or otherwise interfere with OP-1 binding, thereby substantially interfering with osteocalcin induction in the presence of a limiting amount of OP-1.

- 41 -

Mineralization is determined on long term cultures (13 day) using a modified von Kossa staining technique on fixed cell layers: cells are fixed in fresh 4% paraformaldehyde at 23°C for 10 min, following rinsing cold 0.9% NaCl. Fixed cells then are stained for endogenous alkaline phosphatase at pH 9.5 for 10 min, using a commercially available kit (Sigma, Inc.) Purple stained cells then are dehydrated with methanol and air dried. After 30 min incubation in 3% AgNO<sub>3</sub> in the dark, H<sub>2</sub>O-rinsed samples are exposed for 30 sec to 254 nm UV light to develop the black silver-stained phosphate nodules. Individual mineralized foci (at least 20 mm in size) are counted under a dissecting microscope and expressed as nodules/culture. OP-1 induces a 20-fold increase in initial mineralization rate. Agonists are expected to have similar induction effects; antagonists are expected to inhibit or otherwise interfere with OP-1 binding, thereby inhibiting mineralization induction in the presence of a limiting amount of OP-1.

## 9.2 Induction of a Constructed Reporter Gene

Alternatively, a reporter gene construct can be used to determine the ability of candidate molecule to induce signal transduction across a membrane following receptor binding. For example, PAI-1 protein, (Plasminogen Activator Inhibitor-1) expression can be induced by OP-1 in Mv1Lu- cells (see above). Also, as demonstrated above, ALK-1, when overexpressed in a chemically mutagenized derivative of these cells, can mediate PAI-1 induction in the presence of OP-1.

Accordingly, PAI-1 promoter elements can be fused to a reporter gene and induction of the reporter gene monitored following incubation of the transfected cell with a candidate analog. As one example, the luciferase reporter gene can be used, in, for example, the construct p3TP-Lux described by Wrana *et al.* (1992) *Cell* 71: 1003-1014 and Attisano *et al.* (1993) *Cell* 74: 671-680. This reporter gene construct includes a region of the human PAI-1 gene promoter in combination with three sets of tetradecanoyl phorbol - acetate responsive elements upstream of the luciferase open reading frame.

In a typical assay, transfected cells starved in DMEM containing 0.1% fetal bovine serum and antibiotics (*e.g.*, 100 units/ml penicillin and 50 µg/ml streptomycin) for 6 hrs., and then exposed to ligand for 24 hr. Luciferase activity in the cell lysate then is measured using a luminometer in the luciferase assay system, according to the manufacturer's protocol (Promega).

In Mv1Lu mutant cells, "R mutant" cells co-transfected with ALK-2 and ActRII, OP-1-mediated induction of luciferase activity.

### 9.3 Inhibition of Epithelial cell proliferation

OP-1 is known to inhibit epithelial cells. Thus, the ability of a candidate analog to inhibit  
5 cell proliferation, as measured by  $^3\text{H}$ -thymidine uptake by an epithelial cell can be used in an assay to evaluate signal transduction activity of the candidate. Analogs competent to inhibit epithelial cell growth are contemplated to have particular utility in therapeutic applications where limitation of a proliferating cell population is desired. Such applications include chemotherapies and radiation therapies where limiting the growth of a normally proliferating population of cells can  
10 protect these cells from the cytotoxic effects of these cancer therapies. (see *e.g.*, WO 94/06420). In addition, psoriasis and other tissue disorders resulting from uncontrolled cell proliferation, including benign and malignant neoplasies, can be modulated by use of an OP-1 analog.

As an example, mink lung epithelial cell growth is inhibited by OP-1. (see, PCT US93/08885; WO 94/06420.) As described above, derivatives of these cells [*e.g.*, "R-4 mutants",  
15 clone 4-2, Laiho *et al.* (1990) *J. Biol. Chem.* 265: 18518-18524] can be transfected with DNA encoding OP-1-specific receptors and induced to express these receptors. The transfected cells, then can be assayed for a candidate analog's ability to block cell growth. As one example, when R-4 cells are transfected with ALK-1 under a  $\text{Zn}^{2+}$ -inducible promoter, and induced to express the receptor following induction with  $\text{ZnCl}_2$ , cell growth can be inhibited in the presence of OP-1 in a  
20 dose dependent manner.

In a typical assay, cells are seeded in 24-well cell culture plates at a density of 104 cells per well in DMEM with 10% FBS, and incubated overnight. The medium is replaced with DMEM containing 0.2% FBS and 100  $\mu\text{M}$   $\text{ZnCl}_2$ , and the cells are incubated for 5 hrs, after which the medium is replaced with fresh DMEM containing 0.2% FBS, 100  $\mu\text{M}$   $\text{ZnCl}_2$  and  
25 various concentrations of OP-1 or an analog candidate. After 16 h of incubation, 0.25 Ci of  $^3\text{H}$ -thymidine (Amersham) are added and the cells incubated for an additional 2 hrs. Thereafter, the cells are fixed in 10% trichloroacetic acid for more than 15 min on ice, and solubilized with 1 M NaOH. The cell extracts are neutralized with 1 M HCl and  $^3\text{H}$  radioactivity determined in a liquid scintillation counter.

### Example 10. SCREENING ASSAY FOR COMPOUNDS WHICH ALTER ENDOGENOUS OP-1 RECEPTOR EXPRESSION LEVELS

Candidate compound(s) which can be administered to affect the level of a given endogenous OP-1 receptor can be found using the following screening assay, in which the level of OP-1 receptor production by a cell type which produces measurable levels of the receptor is determined by incubating the cell in culture with and without the candidate compound, in order to assess the effects of the compound on the cell. This can also be accomplished by detection of the OP-1 receptor either at the protein level by Western blot or immunolocalization, or at the RNA level by Northern blot or *in situ* hybridization. The protocol is based on a procedure for identifying compounds which alter endogenous levels of OP-1 expression, a detailed description also may be found in PCT US 92/07359, incorporated herein by reference.

Cell cultures of, for example, bone, brain, intestine, lung, heart, eye, breast, gonads, kidney, adrenals, urinary bladder, brain, or other organs, may be prepared as described widely in the literature. For example, kidneys may be explanted from neonatal or new born or young or adult rodents (mouse or rat) and used in organ culture as whole or sliced (1-4 mm) tissues. Primary tissue cultures and established cell lines, also derived from kidney, adrenals, urinary, bladder, brain, mammary, or other tissues may be established in multiwell plates (6 well or 24 well) according to conventional cell culture techniques, and are cultured in the absence or presence of serum for a period of time (1-7 days). Cells can be cultured, for example, in Dulbecco's Modified Eagle medium (Gibco, Long Island, NY) containing serum (*e.g.*, fetal calf serum at 1%-10%, Gibco) or in serum-deprived medium, as desired, or in defined medium (*e.g.*, containing insulin, transferring, glucose, albumin, or other growth factors).

Cell samples for testing the level of OP-1 receptor production are collected periodically and evaluated for receptor production by immunoblot analysis (Sambrook *et al.*, eds., 1989, *Molecular Cloning*, Cold Spring Harbor Press, Cold Spring Harbor, NY), or, alternatively, a portion of the cell culture itself can be collected periodically and used to prepare poly(A)<sup>+</sup> RNA for mRNA analysis by Northern blot analysis. To monitor *de novo* receptor synthesis, some cultures are labeled according to conventional procedures with an <sup>35</sup>S-methionine/<sup>35</sup>S-cysteine mixture for 6-24 hours and then evaluated to quantitate receptor synthesis by conventional immunoassay methods. Alternatively, anti-receptor antibodies may be labeled and incubated with the cells or cell lysates, and the bound complexes detected and quantitated by conventional means,

such as those described hereinabove. Northern blots may be performed using a portion of the OP-1 receptor coding sequence to create hybridization probes, and following the RNA hybridization protocol described herein.

#### Example 11. GENERAL FORMULATION/ ADMINISTRATION CONSIDERATIONS

5       The analogs and constructs described herein can be provided to an individual as part of a therapy to enhance, inhibit, or otherwise modulate the *in vivo* binding interaction between OP-1 and one or more OP-1-specific cell surface receptors. The molecules then comprise part of a pharmaceutical composition as described herein below and can be administered by any suitable means, preferably directly or systemically, *e.g.*, parenterally or orally. Where the therapeutic  
10   molecule is to be provided directly (*e.g.*, locally, as by injection, to a desired tissue site), or parenterally, such as by intravenous, subcutaneous, intramuscular, intraorbital, ophthalmic, intraventricular, intracranial, intracapsular, intraspinal, intracisternal, intraperitoneal, buccal, rectal, vaginal, intranasal or by aerosol administration, the therapeutic preferably comprises part of an aqueous solution. The solution preferably is physiologically acceptable so that in addition to  
15   delivery of the desired morphogen to the patient, the solution does not otherwise adversely affect the patient's electrolyte and volume balance. The aqueous medium for the therapeutic molecule thus may comprise normal physiologic saline (0.9% NaCl, 0.15 M), pH 7-7.4 or other pharmaceutically acceptable salts thereof.

Useful solutions for oral or parenteral administration can be prepared by any of the  
20   methods well known in the pharmaceutical art, described, for example, in *Remington's Pharmaceutical Sciences*, (Gennaro, A., ed.), Mack Pub., 1990. Formulations may include, for example, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes, and the like. Formulations for direct administration, in particular, can include glycerol and other compositions of high viscosity. Biocompatible, preferably bioresorbable  
25   polymers, including, for example, hyaluronic acid, collagen, tricalcium phosphate, polybutyrate, polylactide, polyglycolide and lactide/glycolide copolymers, may be useful excipients to control the release of the morphogen *in vivo*.

Other potentially useful parenteral delivery systems for these therapeutic molecules include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and  
30   liposomes. Formulations for inhalation administration may contain as excipients, for example, lactose, or can be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether,

- 45 -

glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally.

Alternatively, the morphogens described herein may be administered orally.

The therapeutic molecules can also be associated with means for targeting the therapeutic to a desired tissue. For example, tetracycline and diphosphonates (bisphosphonates) are known to bind to bone mineral, particularly at zones of bone remodeling, when they are provided systemically in a mammal. Accordingly, these molecules may be included as useful agents for targeting therapeutics to bone tissue. Alternatively, an antibody or other binding protein that interacts specifically with a surface molecule on the desired target tissue cells can also be used.

Such targeting molecules further can be covalently associated to the therapeutic molecule *e.g.*, by chemical crosslinking, or by using standard genetic engineering means to create, for example, an acid labile bond such as an Asp-Pro linkage. Useful targeting molecules can be designed, for example, using the single chain binding site technology disclosed, for example, in U.S. Pat. No. 5,091,513.

Finally, therapeutic molecules can be administered alone or in combination with other molecules known to have a beneficial effect on tissue morphogenesis, including molecules capable of tissue repair and regeneration and/or inhibiting inflammation. Examples of useful cofactors for stimulating bone tissue growth in osteoporotic individuals, for example, include but are not limited to, vitamin D<sub>3</sub>, calcitonin, prostaglandins, parathyroid hormone, dexamethasone, estrogen and IGF-I or IGF-II. Useful cofactors for nerve tissue repair and regeneration can include nerve growth factors. Other useful cofactors include symptom-alleviating cofactors, including antiseptics, antibiotics, antiviral and antifungal agents and analgesics and anesthetics.

Therapeutic molecules further can be formulated into pharmaceutical compositions by admixture with pharmaceutically acceptable nontoxic excipients and carriers. As noted above, such compositions can be prepared for parenteral administration, particularly in the form of liquid solutions or suspensions; for oral administration, particularly in the form of tablets or capsules; or intranasally, particularly in the form of powders, nasal drops or aerosols. Where adhesion to a tissue surface is desired the composition may include the morphogen dispersed in a fibrinogen-thrombin composition or other bioadhesive such as is disclosed, for example in PCT US91/09275,

the disclosure of which is incorporated herein by reference. The composition then can be painted, sprayed or otherwise applied to the desired tissue surface.

The compositions can be formulated for parenteral or oral administration to humans or other mammals in therapeutically effective amounts, *e.g.*, amounts which provide appropriate concentrations of the analog to target tissue for a time sufficient to induce the desired effect.

Where the analog is to be used as part of a transplant procedure, it can be provided to the living tissue or organ to be transplanted prior to removal of tissue or organ from the donor. The analog may be provided to the donor host directly, as by injection of a formulation comprising the analog into the tissue, or indirectly, *e.g.*, by oral or parenteral administration, using any of the means described above.

Alternatively or, in addition, once removed from the donor, the organ or living tissue can be placed in a preservation solution containing the therapeutic molecule. In addition, the recipient also preferably is provided with the analog just prior to, or concomitant with, transplantation. In all cases, the analog can be administered directly to the tissue at risk, as by injection to the tissue, or it may be provided systemically, either by oral or parenteral administration, using any of the methods and formulations described herein and/or known in the art.

Where the therapeutic molecule comprises part of a tissue or organ preservation solution, any commercially available preservation solution can be used to advantage. For example, useful solutions known in the art include Collins solution, Wisconsin solution, Belzer solution, Eurocollins solution and lactated Ringer's solution. Generally, an organ preservation solution usually possesses one or more of the following properties: (a) an osmotic pressure substantially equal to that of the inside of a mammalian cell, (solutions typically are hyperosmolar and have  $K^+$  and/or  $Mg^{2+}$  ions present in an amount sufficient to produce an osmotic pressure slightly higher than the inside of a mammalian cell); (b) the solution typically is capable of maintaining substantially normal ATP levels in the cells; and (c) the solution usually allows optimum maintenance of glucose metabolism in the cells. Organ preservation solutions also may contain anticoagulants, energy sources such as glucose, fructose and other sugars, metabolites, heavy metal chelators, glycerol and other materials of high viscosity to enhance survival at low temperatures, free oxygen radical inhibiting and/or scavenging agents and a pH indicator. A detailed description of preservation solutions and useful components can be found, for example, in US Patent No. 5,002,965, the disclosure of which is incorporated herein by reference.



- 47 -

As will be appreciated by those skilled in the art, the concentration of the compounds described in a therapeutic composition will vary depending upon a number of factors, including the dosage of the drug to be administered, the chemical characteristics (*e.g.*, hydrophobicity) of the compounds employed, and the route of administration. The preferred dosage of drug to be administered also is likely to depend on such variables as the type and extent of tissue loss or defect, the overall health status of the particular patient, the relative biological efficacy of the compound selected, the formulation of the compound, the presence and types of excipients in the formulation, and the route of administration. In general terms, the therapeutic molecules of this invention may be provided to an individual where typical dose ranges are from about 10 ng/kg to about 1 g/kg of body weight per day; a preferred dose range being from about 0.1 mg/kg to 100 mg/kg of body weight. No obvious morphogen-induced pathological lesions are induced when mature morphogen (*e.g.*, OP-1, 20 mg) is administered daily to normal growing rats for 21 consecutive days. Moreover, 10 mg systemic injections of morphogen (*e.g.*, OP-1) injected daily for 10 days into normal newborn mice does not produce any gross abnormalities.

## 15 Other Embodiments

The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The present embodiments are therefore to be considered in all respects as illustrative and not restrictive, the scope of the invention being indicated by the appended claims rather than by the foregoing description, and all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

- 48 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: ICHIJO, HIDENORI  
NISHITOH, HIDEKI  
SAMPATH, KUBER T.
- (ii) TITLE OF INVENTION: NOVEL SIGNALLING RECEPTOR FOR MORPHOGENIC PROTEINS
- (iii) NUMBER OF SEQUENCES: 5
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Testa, Hurwitz & Thibault
  - (B) STREET: 125 High St.
  - (C) CITY: Boston
  - (D) STATE: MA
  - (E) COUNTRY: USA
  - (F) ZIP: 02110
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: KELLEY Esq., Robin D.
  - (B) REGISTRATION NUMBER: 34,637
  - (C) REFERENCE/DOCKET NUMBER: CRP-117
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (617) 248-7000
  - (B) TELEFAX: (617) 248-7100

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1509 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 1..1509
  - (D) OTHER INFORMATION: /product= "Human ALK-1"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG ACC TTG GGC TCC CCC AGG AAA GGC CTT CTG ATG CTG CTG ATG GCC  
 Met Thr Leu Gly Ser Pro Arg Lys Gly Leu Leu Met Leu Leu Met Ala  
 1 5 10 15

48

- 49 -

TTG	GTG	ACC	CAG	GGA	GAC	CCT	GTG	AAG	CCG	TCT	CGG	GGC	CCG	CTG	GTG	96
Leu	Val	Thr	Gln	Gly	Asp	Pro	Val	Lys	Pro	Ser	Arg	Gly	Pro	Leu	Val	
			20					25					30			
ACC	TGC	ACG	TGT	GAG	AGC	CCA	CAT	TGC	AAG	GGG	CCT	ACC	TGC	CGG	GGG	144
Thr	Cys	Thr	Cys	Glu	Ser	Pro	His	Cys	Lys	Gly	Pro	Thr	Cys	Arg	Gly	
		35					40					45				
GCC	TGG	TGC	ACA	GTA	GTG	CTG	GTG	CGG	GAG	GAG	GGG	AGG	CAC	CCC	CAG	192
Ala	Trp	Cys	Thr	Val	Val	Leu	Val	Arg	Glu	Glu	Gly	Arg	His	Pro	Gln	
	50					55					60					
GAA	CAT	CGG	GGC	TGC	GGG	AAC	TTG	CAC	AGG	GAG	CTC	TGC	AGG	GGG	CGC	240
Glu	His	Arg	Gly	Cys	Gly	Asn	Leu	His	Arg	Glu	Leu	Cys	Arg	Gly	Arg	
	65				70					75					80	
CCC	ACC	GAG	TTC	GTC	AAC	CAC	TAC	TGC	TGC	GAC	AGC	CAC	CTC	TGC	AAC	288
Pro	Thr	Glu	Phe	Val	Asn	His	Tyr	Cys	Cys	Asp	Ser	His	Leu	Cys	Asn	
			85					90						95		
CAC	AAC	GTG	TCC	CTG	GTG	CTG	GAG	GCC	ACC	CAA	CCT	CCT	TCG	GAG	CAG	336
His	Asn	Val	Ser	Leu	Val	Leu	Glu	Ala	Thr	Gln	Pro	Pro	Ser	Glu	Gln	
			100					105					110			
CCG	GGA	ACA	GAT	GGC	CAG	CTG	GCC	CTG	ATC	CTG	GGC	CCC	GTG	CTG	GCC	384
Pro	Gly	Thr	Asp	Gly	Gln	Leu	Ala	Leu	Ile	Leu	Gly	Pro	Val	Leu	Ala	
		115					120					125				
TTG	CTG	GCC	CTG	GTG	GCC	CTG	GGT	GTC	CTG	GGC	CTG	TGG	CAT	GTC	CGA	432
Leu	Leu	Ala	Leu	Val	Ala	Leu	Gly	Val	Leu	Gly	Leu	Trp	His	Val	Arg	
	130					135					140					
CGG	AGG	CAG	GAG	AAG	CAG	CGT	GGC	CTG	CAC	AGC	GAG	CTG	GGA	GAG	TCC	480
Arg	Arg	Gln	Glu	Lys	Gln	Arg	Gly	Leu	His	Ser	Glu	Leu	Gly	Glu	Ser	
	145				150					155					160	
AGT	CTC	ATC	CTG	AAA	GCA	TCT	GAG	CAG	GGC	GAC	ACG	ATG	TTG	GGG	GAC	528
Ser	Leu	Ile	Leu	Lys	Ala	Ser	Glu	Gln	Gly	Asp	Thr	Met	Leu	Gly	Asp	
				165					170					175		
CTC	CTG	GAC	AGT	GAC	TGC	ACC	ACA	GGG	AGT	GGC	TCA	GGG	CTC	CCC	TTC	576
Leu	Leu	Asp	Ser	Asp	Cys	Thr	Thr	Gly	Ser	Gly	Ser	Gly	Leu	Pro	Phe	
			180					185					190			
CTG	GTG	CAG	AGG	ACA	GTG	GCA	CGG	CAG	GTT	GCC	TTG	GTG	GAG	TGT	GTG	624
Leu	Val	Gln	Arg	Thr	Val	Ala	Arg	Gln	Val	Ala	Leu	Val	Glu	Cys	Val	
		195					200					205				
GGA	AAA	GGC	CGC	TAT	GGC	GAA	GTG	TGG	CGG	GGC	TTG	TGG	CAC	GGT	GAG	672
Gly	Lys	Gly	Arg	Tyr	Gly	Glu	Val	Trp	Arg	Gly	Leu	Trp	His	Gly	Glu	
	210					215					220					
AGT	GTG	GCC	GTC	AAG	ATC	TTC	TCC	TCG	AGG	GAT	GAA	CAG	TCC	TGG	TTC	720
Ser	Val	Ala	Val	Lys	Ile	Phe	Ser	Ser	Arg	Asp	Glu	Gln	Ser	Trp	Phe	
	225				230					235					240	
CGG	GAG	ACT	GAG	ATC	TAT	AAC	ACA	GTA	TTG	CTC	AGA	CAC	GAC	AAC	ATC	768
Arg	Glu	Thr	Glu	Ile	Tyr	Asn	Thr	Val	Leu	Leu	Arg	His	Asp	Asn	Ile	
				245					250					255		
CTA	GGC	TTC	ATC	GCC	TCA	GAC	ATG	ACC	TCC	CGC	AAC	TCG	AGC	ACG	CAG	816
Leu	Gly	Phe	Ile	Ala	Ser	Asp	Met	Thr	Ser	Arg	Asn	Ser	Ser	Thr	Gln	
			260					265					270			

- 50 -

CTG	TGG	CTC	ATC	ACG	CAC	TAC	CAC	GAG	CAC	GGC	TCC	CTC	TAC	GAC	TTT	864
Leu	Trp	Leu	Ile	Thr	His	Tyr	His	Glu	His	Gly	Ser	Leu	Tyr	Asp	Phe	
		275					280					285				
CTG	CAG	AGA	CAG	ACG	CTG	GAG	CCC	CAT	CTG	GCT	CTG	AGG	CTA	GCT	GTG	912
Leu	Gln	Arg	Gln	Thr	Leu	Glu	Pro	His	Leu	Ala	Leu	Arg	Leu	Ala	Val	
	290					295					300					
TCC	GCG	GCA	TGC	GGC	CTG	GCG	CAC	CTG	CAC	GTG	GAG	ATC	TTC	GGT	ACA	960
Ser	Ala	Ala	Cys	Gly	Leu	Ala	His	Leu	His	Val	Glu	Ile	Phe	Gly	Thr	
305					310					315					320	
CAG	GGC	AAA	CCA	GCC	ATT	GCC	CAC	CGC	GAC	TTC	AAG	AGC	CGC	AAT	GTG	1008
Gln	Gly	Lys	Pro	Ala	Ile	Ala	His	Arg	Asp	Phe	Lys	Ser	Arg	Asn	Val	
				325					330					335		
CTG	GTC	AAG	AGC	AAC	CTG	CAG	TGT	TGC	ATC	GCC	GAC	CTG	GGC	CTG	GCT	1056
Leu	Val	Lys	Ser	Asn	Leu	Gln	Cys	Cys	Ile	Ala	Asp	Leu	Gly	Leu	Ala	
			340					345					350			
GTG	ATG	CAC	TCA	CAG	GGC	AGC	GAT	TAC	CTG	GAC	ATC	GGC	AAC	AAC	CCG	1104
Val	Met	His	Ser	Gln	Gly	Ser	Asp	Tyr	Leu	Asp	Ile	Gly	Asn	Asn	Pro	
		355					360					365				
AGA	GTG	GGC	ACC	AAG	CGG	TAC	ATG	GCA	CCC	GAG	GTG	CTG	GAC	GAG	CAG	1152
Arg	Val	Gly	Thr	Lys	Arg	Tyr	Met	Ala	Pro	Glu	Val	Leu	Asp	Glu	Gln	
	370					375					380					
ATC	CGC	ACG	GAC	TGC	TTT	GAG	TCC	TAC	AAG	TGG	ACT	GAC	ATC	TGG	GCC	1200
Ile	Arg	Thr	Asp	Cys	Phe	Glu	Ser	Tyr	Lys	Trp	Thr	Asp	Ile	Trp	Ala	
385					390					395					400	
TTT	GGC	CTG	GTG	CTG	TGG	GAG	ATT	GCC	CGC	CGG	ACC	ATC	GTG	AAT	GGC	1248
Phe	Gly	Leu	Val	Leu	Trp	Glu	Ile	Ala	Arg	Arg	Thr	Ile	Val	Asn	Gly	
				405				410						415		
ATC	GTG	GAG	GAC	TAT	AGA	CCA	CCC	TTC	TAT	GAT	GTG	GTG	CCC	AAT	GAC	1296
Ile	Val	Glu	Asp	Tyr	Arg	Pro	Pro	Phe	Tyr	Asp	Val	Val	Pro	Asn	Asp	
			420					425					430			
CCC	AGC	TTT	GAG	GAC	ATG	AAG	AAG	GTG	GTG	TGT	GTG	GAT	CAG	CAG	ACC	1344
Pro	Ser	Phe	Glu	Asp	Met	Lys	Lys	Val	Val	Cys	Val	Asp	Gln	Gln	Thr	
		435					440					445				
CCC	ACC	ATC	CCT	AAC	CGG	CTG	GCT	GCA	GAC	CCG	GTC	CTC	TCA	GGC	CTA	1392
Pro	Thr	Ile	Pro	Asn	Arg	Leu	Ala	Ala	Asp	Pro	Val	Leu	Ser	Gly	Leu	
		450				455					460					
GCT	CAG	ATG	ATG	CGG	GAG	TGC	TGG	TAC	CCA	AAC	CCC	TCT	GCC	CGA	CTC	1440
Ala	Gln	Met	Met	Arg	Glu	Cys	Trp	Tyr	Pro	Asn	Pro	Ser	Ala	Arg	Leu	
465					470					475					480	
ACC	GCG	CTG	CGG	ATC	AAG	AAG	ACA	CTA	CAA	AAA	ATT	AGC	AAC	AGT	CCA	1488
Thr	Ala	Leu	Arg	Ile	Lys	Lys	Thr	Leu	Gln	Lys	Ile	Ser	Asn	Ser	Pro	
				485				490						495		
GAG	AAG	CCT	AAA	GTG	ATT	CAA										1509
Glu	Lys	Pro	Lys	Val	Ile	Gln										
				500												

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 503 amino acids

- 51 -

(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Thr Leu Gly Ser Pro Arg Lys Gly Leu Leu Met Leu Leu Met Ala
 1      5      10      15
Leu Val Thr Gln Gly Asp Pro Val Lys Pro Ser Arg Gly Pro Leu Val
 20      25      30
Thr Cys Thr Cys Glu Ser Pro His Cys Lys Gly Pro Thr Cys Arg Gly
 35      40      45
Ala Trp Cys Thr Val Val Leu Val Arg Glu Glu Gly Arg His Pro Gln
 50      55      60
Glu His Arg Gly Cys Gly Asn Leu His Arg Glu Leu Cys Arg Gly Arg
 65      70      75      80
Pro Thr Glu Phe Val Asn His Tyr Cys Cys Asp Ser His Leu Cys Asn
 85      90      95
His Asn Val Ser Leu Val Leu Glu Ala Thr Gln Pro Pro Ser Glu Gln
 100      105      110
Pro Gly Thr Asp Gly Gln Leu Ala Leu Ile Leu Gly Pro Val Leu Ala
 115      120      125
Leu Leu Ala Leu Val Ala Leu Gly Val Leu Gly Leu Trp His Val Arg
 130      135      140
Arg Arg Gln Glu Lys Gln Arg Gly Leu His Ser Glu Leu Gly Glu Ser
 145      150      155      160
Ser Leu Ile Leu Lys Ala Ser Glu Gln Gly Asp Thr Met Leu Gly Asp
 165      170      175
Leu Leu Asp Ser Asp Cys Thr Thr Gly Ser Gly Ser Gly Leu Pro Phe
 180      185      190
Leu Val Gln Arg Thr Val Ala Arg Gln Val Ala Leu Val Glu Cys Val
 195      200      205
Gly Lys Gly Arg Tyr Gly Glu Val Trp Arg Gly Leu Trp His Gly Glu
 210      215      220
Ser Val Ala Val Lys Ile Phe Ser Ser Arg Asp Glu Gln Ser Trp Phe
 225      230      235      240
Arg Glu Thr Glu Ile Tyr Asn Thr Val Leu Leu Arg His Asp Asn Ile
 245      250      255
Leu Gly Phe Ile Ala Ser Asp Met Thr Ser Arg Asn Ser Ser Thr Gln
 260      265      270
Leu Trp Leu Ile Thr His Tyr His Glu His Gly Ser Leu Tyr Asp Phe
 275      280      285
Leu Gln Arg Gln Thr Leu Glu Pro His Leu Ala Leu Arg Leu Ala Val
 290      295      300
Ser Ala Ala Cys Gly Leu Ala His Leu His Val Glu Ile Phe Gly Thr

```

- 52 -

```

305          310          315          320
Gln Gly Lys Pro Ala Ile Ala His Arg Asp Phe Lys Ser Arg Asn Val
          325          330          335
Leu Val Lys Ser Asn Leu Gln Cys Cys Ile Ala Asp Leu Gly Leu Ala
          340          345          350
Val Met His Ser Gln Gly Ser Asp Tyr Leu Asp Ile Gly Asn Asn Pro
          355          360          365
Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Gln
          370          375          380
Ile Arg Thr Asp Cys Phe Glu Ser Tyr Lys Trp Thr Asp Ile Trp Ala
          385          390          395
Phe Gly Leu Val Leu Trp Glu Ile Ala Arg Arg Thr Ile Val Asn Gly
          405          410          415
Ile Val Glu Asp Tyr Arg Pro Pro Phe Tyr Asp Val Val Pro Asn Asp
          420          425          430
Pro Ser Phe Glu Asp Met Lys Lys Val Val Cys Val Asp Gln Gln Thr
          435          440          445
Pro Thr Ile Pro Asn Arg Leu Ala Ala Asp Pro Val Leu Ser Gly Leu
          450          455          460
Ala Gln Met Met Arg Glu Cys Trp Tyr Pro Asn Pro Ser Ala Arg Leu
          465          470          475
Thr Ala Leu Arg Ile Lys Lys Thr Leu Gln Lys Ile Ser Asn Ser Pro
          485          490          495
Glu Lys Pro Lys Val Ile Gln
          500

```

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1822 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
  - (B) LOCATION: 49..1341
  - (D) OTHER INFORMATION: /function= "Osteogenic Protein"
- /product= "OP-1"
- /standard\_name= "OP-1"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

GGTGCGGGCC CGGAGCCCGG AGCCCGGGTA GCGCGTAGAG CCGGCGCG ATG CAC GTG
                                         Met His Val
                                         1
CGC TCA CTG CGA GCT GCG GCG CCG CAC AGC TTC GTG GCG CTC TGG GCA
Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala Leu Trp Ala

```

57

105

5					10					15						
CCC Pro 20	CTG Leu	TTC Phe	CTG Leu	CTG Leu	CGC Arg 25	TCC Ser	GCC Ala	CTG Leu	GCC Ala	GAC Asp 30	TTC Phe	AGC Ser	CTG Leu	GAC Asp	AAC Asn 35	153
GAG Glu	GTG Val	CAC His	TCG Ser	AGC Ser 40	TTC Phe	ATC Ile	CAC His	CGG Arg	CGC Arg 45	CTC Leu	CGC Arg	AGC Ser	CAG Gln	GAG Glu 50	CGG Arg	201
CGG Arg	GAG Glu	ATG Met	CAG Gln 55	CGC Arg	GAG Glu	ATC Ile	CTC Leu	TCC Ser 60	ATT Ile	TTG Leu	GGC Gly	TTG Leu	CCC Pro 65	CAC His	CGC Arg	249
CCG Pro	CGC Arg	CCG Pro 70	CAC His	CTC Leu	CAG Gln	GGC Gly	AAG Lys 75	CAC His	AAC Asn	TCG Ser	GCA Ala	CCC Pro 80	ATG Met	TTC Phe	ATG Met	297
CTG Leu	GAC Asp 85	CTG Leu	TAC Tyr	AAC Asn	GCC Ala	ATG Met 90	GCG Ala	GTG Val	GAG Glu	GAG Glu	GGC Gly 95	GGC Gly	GGG Gly	CCC Pro	GGC Gly	345
GGC Gly 100	CAG Gln	GGC Gly	TTC Phe	TCC Ser	TAC Tyr 105	CCC Pro	TAC Tyr	AAG Lys	GCC Ala	GTC Val 110	TTC Phe	AGT Ser	ACC Thr	CAG Gln	GGC Gly 115	393
CCC Pro	CCT Pro	CTG Leu	GCC Ala	AGC Ser 120	CTG Leu	CAA Gln	GAT Asp	AGC Ser	CAT His 125	TTC Phe	CTC Leu	ACC Thr	GAC Asp	GCC Ala 130	GAC Asp	441
ATG Met	GTC Val	ATG Met	AGC Ser 135	TTC Phe	GTC Val	AAC Asn	CTC Leu	GTG Val 140	GAA Glu	CAT His	GAC Asp	AAG Lys	GAA Glu 145	TTC Phe	TTC Phe	489
CAC His	CCA Pro	CGC Arg 150	TAC Tyr	CAC His	CAT His	CGA Arg	GAG Glu 155	TTC Phe	CGG Arg	TTT Phe	GAT Asp	CTT Leu	TCC Ser 160	AAG Lys	ATC Ile	537
CCA Pro	GAA Glu 165	GGG Gly	GAA Glu	GCT Ala	GTC Val	ACG Thr 170	GCA Ala	GCC Ala	GAA Glu	TTC Phe	CGG Arg 175	ATC Ile	TAC Tyr	AAG Lys	GAC Asp	585
TAC Tyr 180	ATC Ile	CGG Arg	GAA Glu	CGC Arg	TTC Phe 185	GAC Asp	AAT Asn	GAG Glu	ACG Thr	TTC Phe 190	CGG Arg	ATC Ile	AGC Ser	GTT Val	TAT Tyr 195	633
CAG Gln	GTG Val	CTC Leu	CAG Gln	GAG Glu 200	CAC His	TTG Leu	GGC Gly	AGG Arg	GAA Glu 205	TCG Ser	GAT Asp	CTC Leu	TTC Phe	CTG Leu 210	CTC Leu	681
GAC Asp	AGC Ser	CGT Arg	ACC Thr 215	CTC Leu	TGG Trp	GCC Ala	TCG Ser	GAG Glu 220	GAG Glu	GGC Gly	TGG Trp	CTG Leu	GTG Val 225	TTT Phe	GAC Asp	729
ATC Ile	ACA Thr 230	GCC Ala	ACC Thr	AGC Ser	AAC Asn	CAC His	TGG Trp 235	GTG Val	GTC Val	AAT Asn	CCG Pro	CGG Arg 240	CAC His	AAC Asn	CTG Leu	777
GGC Gly 245	CTG Leu	CAG Gln	CTC Leu	TCG Ser	GTG Val	GAG Glu 250	ACG Thr	CTG Leu	GAT Asp	GGG Gly	CAG Gln 255	AGC Ser	ATC Ile	AAC Asn	CCC Pro	825
AAG Lys 260	TTG Leu	GCG Ala	GGC Gly	CTG Leu	ATT Ile 265	GGG Gly	CGG Arg	CAC His	GGG Gly	CCC Pro 270	CAG Gln	AAC Asn	AAG Lys	CAG Gln	CCC Pro 275	873

- 54 -

TTC ATG GTG GCT TTC TTC AAG GCC ACG GAG GTC CAC TTC CGC AGC ATC Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe Arg Ser Ile 280 285 290	921
CGG TCC ACG GGG AGC AAA CAG CGC AGC CAG AAC CGC TCC AAG ACG CCC Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser Lys Thr Pro 295 300 305	969
AAG AAC CAG GAA GCC CTG CGG ATG GCC AAC GTG GCA GAG AAC AGC AGC Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu Asn Ser Ser 310 315 320	1017
AGC GAC CAG AGG CAG GCC TGT AAG AAG CAC GAG CTG TAT GTC AGC TTC Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val Ser Phe 325 330 335	1065
CGA GAC CTG GGC TGG CAG GAC TGG ATC ATC GCG CCT GAA GGC TAC GCC Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala 340 345 350 355	1113
GCC TAC TAC TGT GAG GGG GAG TGT GCC TTC CCT CTG AAC TCC TAC ATG Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met 360 365 370	1161
AAC GCC ACC AAC CAC GCC ATC GTG CAG ACG CTG GTC CAC TTC ATC AAC Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn 375 380 385	1209
CCG GAA ACG GTG CCC AAG CCC TGC TGT GCG CCC ACG CAG CTC AAT GCC Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala 390 395 400	1257
ATC TCC GTC CTC TAC TTC GAT GAC AGC TCC AAC GTC ATC CTG AAG AAA Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys 405 410 415	1305
TAC AGA AAC ATG GTG GTC CGG GCC TGT GGC TGC CAC TAGCTCCTCC Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His 420 425 430	1351
GAGAATTCAG ACCCTTTGGG GCCAAGTTTT TCTGGATCCT CCATTGCTCG CTTTGGCCAG	1411
GAACCAGCAG ACCAACTGCC TTTTGTGAGA CCTTCCCCTC CCTATCCCCA ACTTTAAAGG	1471
TGTGAGAGTA TTAGGAAACA TGAGCAGCAT ATGGCTTTTG ATCAGTTTTT CAGTGGCAGC	1531
ATCCAATGAA CAAGATCCTA CAAGCTGTGC AGGCAAAACC TAGCAGGAAA AAAAAACAAC	1591
GCATAAAGAA AAATGGCCGG GCCAGGTCAT TGGCTGGGAA GTCTCAGCCA TGCACGGACT	1651
CGTTTCCAGA GGTAATTATG AGCGCCTACC AGCCAGGCCA CCCAGCCGTG GGAGGAAGGG	1711
GGCGTGGCAA GGGGTGGGCA CATTGGTGTC TGTGCGAAAG GAAAATTGAC CCGGAAGTTC	1771
CTGTAATAAA TGTCACAATA AAACGAATGA ATGAAAAAAA AAAAAAAAAA A	1822

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 431 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear



- 55 -

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala
 1           5           10           15
Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser
          20           25           30
Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser
          35           40           45
Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu
          50           55           60
Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro
 65           70           75           80
Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly
          85           90           95
Gly Pro Gly Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser
          100          105          110
Thr Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr
          115          120          125
Asp Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys
 130          135          140
Glu Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu
 145          150          155          160
Ser Lys Ile Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile
          165          170          175
Tyr Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Arg Ile
          180          185          190
Ser Val Tyr Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu
          195          200          205
Phe Leu Leu Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu
 210          215          220
Val Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg
 225          230          235          240
His Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser
          245          250          255
Ile Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn
          260          265          270
Lys Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe
          275          280          285
Arg Ser Ile Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser
 290          295          300
Lys Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu
 305          310          315          320
Asn Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr

```

325

330

335

Val Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu  
340 345 350

Gly Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn  
355 360 365

Ser Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His  
370 375 380

Phe Ile Asn Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln  
385 390 395 400

Leu Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile  
405 410 415

Leu Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His  
420 425 430

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 102 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..102

(D) OTHER INFORMATION: /label= OPX

/note= "Each Xaa is independently selected from a group of one or more specified amino acids as defined in the Specification."

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Cys Xaa Xaa His Glu Leu Tyr Val Xaa Phe Xaa Asp Leu Gly Trp Xaa  
1 5 10 15

Asp Trp Xaa Ile Ala Pro Xaa Gly Tyr Xaa Ala Tyr Tyr Cys Glu Gly  
20 25 30

Glu Cys Xaa Phe Pro Leu Xaa Ser Xaa Met Asn Ala Thr Asn His Ala  
35 40 45

Ile Xaa Gln Xaa Leu Val His Xaa Xaa Xaa Pro Xaa Xaa Val Pro Lys  
50 55 60

Xaa Cys Cys Ala Pro Thr Xaa Leu Xaa Ala Xaa Ser Val Leu Tyr Xaa  
65 70 75 80

Asp Xaa Ser Xaa Asn Val Xaa Leu Xaa Lys Xaa Arg Asn Met Val Val  
85 90 95

Xaa Ala Cys Gly Cys His  
100

- 57 -

What is claimed is:

- 1 1. An isolated analog of OP-1 which binds to the ligand binding domain of a cell surface  
2 receptor defined by the amino acid sequence of Seq. ID No. 2 (ALK-1) or a sequence  
3 variant thereof,  
4 said analog having greater than 60% homology with the C-terminal 96 amino acids of the  
5 sequence defined by Seq. ID No. 4 (OP-1).
- 1 2. The analog of claim 1 wherein the analog shares greater than 60% identity with the  
2 C-terminal 96 amino acids of the sequence defined by Seq. ID No. 4 (OP-1).
- 1 3. The analog of claim 1 wherein said analog has specific binding affinity for a Type II cell  
2 surface receptor.
- 1 4. The analog of claim 3 wherein said Type II receptor has binding affinity for activin or  
2 BMP-4.
- 1 5. An isolated analog of OP-1 which binds to the ligand binding domain of a cell  
2 surface receptor defined by the amino acid sequence of Seq. ID No. 2 (ALK-1) or  
3 a sequence variant thereof, said analog defined by OPX, Seq. ID No. 5.
- 1 6. A method for identifying an OP-1 analog, said analog being characterized as  
2 having substantially the same binding affinity for a cell surface receptor as OP-1,  
3 the method comprising the steps of:
  - 4 (a) providing a sample containing a protein consisting of a polypeptide chain  
5 comprising an amino acid sequence defined Seq. ID No. 2 (ALK-1), or an  
6 OP-1-specific receptor analog thereof;
  - 7 (b) contacting said sample with a candidate OP-1 analog; and

- 58 -

8 (c) detecting specific binding between said candidate OP-1 analog and said protein of  
9 step (a).

1 7. A method for identifying an OP-1 analog, said analog being characterized as  
2 having substantially the same binding affinity for a cell surface receptor as OP-1,  
3 the method comprising the steps of:

4 (a) providing a sample containing a protein consisting of a polypeptide chain  
5 comprising an amino acid sequence defined Seq. ID No. 2 (ALK-1), or an  
6 OP-1-specific receptor analog thereof;

7 (b) contacting said sample with a candidate OP-1 analog; and

8 (c) detecting induction of an OP-1-mediated cellular response.

1 8. The method of claim 6 or 7 wherein said sample further comprises part or all of a Type II  
2 serine/threonine kinase receptor protein having binding affinity for OP-1, activin or  
3 BMP-4.

1 9. A method for producing an OP-1 analog, the method comprising the steps of:

2 (a) obtaining, by the method of claim 6 or 7, a candidate OP-1 analog; and

3 (b) producing either said candidate analog or a second OP-1 analog derived from said  
4 candidate and having substantially the same OP-1 receptor-binding domain as said  
5 candidate.

1 10. A kit for identifying OP-1 or a candidate OP-1 analog in a sample, the kit comprising:

2 (a) a receptacle adapted to receive a sample and containing a protein consisting of a  
3 polypeptide chain comprising an amino acid sequence defined Seq. ID No. 2  
4 (ALK-1), or an OP-1-specific receptor analog thereof;

- 59 -

5 (b) means for detecting interaction of OP-1 or a candidate OP-1 analog with said protein of  
6 part (a), said OP-1 or candidate analog comprising part of said sample provided to said  
7 receptacle.

1 11. The kit of claim 10 wherein said means in part (b) comprises either

2 (i) means for detecting specific binding interaction of OP-1 or said candidate  
3 analog with said protein; or

4 (ii) means for detecting induction of an OP-1-mediated cellular response.

1 12. The kit of claim 10 or 11 further comprising a serine/threonine Type II receptor having  
2 binding specificity for OP-1, activin or BMP-4.

1 13. An isolated ligand-receptor complex comprising two molecules, the first said molecule  
2 defining said ligand and comprising at least the C-terminal 96 amino acids of OP-1  
3 (residues 335-431 of Seq. ID No. 4) or an analog thereof, and the second said molecule  
4 defining said receptor and consisting of a polypeptide chain comprising an amino acid  
5 sequence defined Seq. ID No. 2 (ALK-1), or an OP-1-specific receptor analog thereof.

1 14. The complex of claim 13 further comprising part or all of a Type II serine/threonine  
2 kinase receptor.

1 15. An isolated binding partner having specific binding affinity for an epitope on a ligand-  
2 receptor complex, said complex being characterized as comprising an OP-1 protein or an  
3 analog thereof in specific binding interaction with the ligand binding domain of a cell  
4 surface receptor defined by Seq. ID No. 2 (ALK-1), or an OP-1-specific receptor analog  
5 thereof, said binding partner having substantially no binding affinity for the uncomplexed  
6 form of said OP-1 protein or OP-1 analog.

- 60 -

- 1 16. The isolated binding partner of claim 15 wherein said binding partner is further  
2 characterized as having substantially no binding affinity for the uncomplexed form of said  
3 cell surface receptor protein or said analog thereof.
- 1 17. A method for
- 2 (i) antagonizing OP-1 binding to a cell surface receptor; or
- 3 (ii) antagonizing induction of an OP-1 mediated cellular response,  
4 the method comprising the step of providing the OP-1 analog of claim 1 or  
5 2 to an OP-1-responsive cell.
- 1 18. A method for antagonizing
- 2 (i) OP-1 binding to a cell surface receptor; or
- 3 (ii) induction of an OP-1 mediated cellular response, the method comprising  
4 the step of providing to an OP-1-responsive cell a protein consisting of a  
5 polypeptide chain comprising an amino acid sequence defined Seq. ID  
6 No. 2 (ALK-1), or an OP-1-specific receptor analog thereof.
- 1 19. A method for antagonizing activin binding to a cell surface receptor, the method  
2 comprising the step of:  
3 providing a protein having binding specificity for the amino acid sequence defined by Seq.  
4 ID No. 2 (ALK-1) or an OP-1-specific receptor variant thereof, said protein sharing at  
5 least 60% amino acid sequence identity with residue 335-431 of the sequence defined by  
6 Seq. ID No. 4 (OP-1), such that said protein, when provided to a cell expressing said cell  
7 surface receptor, is competent to interact specifically with said receptor, thereby  
8 substantially inhibiting activin binding to said receptor.

- 61 -

1 20. A method for antagonizing BMP-4 binding to a cell surface receptor, the method  
2 comprising the step of:  
3 providing a protein having binding specificity for the ligand binding domain defined by  
4 Seq. ID No. 2 (ALK-1), or an OP-1-specific receptor variant thereof, said protein sharing  
5 at least 60% amino acid sequence identity with residues 335-431 of the sequence defined  
6 by Seq. ID No. 4 (OP-1),  
7 such that said protein, when provided to a cell expressing a said cell surface receptor with  
8 said cell, is competent to interact specifically with said receptor, thereby substantially  
9 inhibiting BMP-4 binding to said receptor.

1/2


cDNA	ALK-1		ALK-3		ALK-5	
	-	+	-	+	-	+
ZnCl <sub>2</sub>	-	+	-	+	-	+
ligand	-	O - O β A	-	O - O β A	-	O - O β A
PAI-1 →						

Fig. 1

SUBSTITUTE SHEET (RULE 26)



2/2




Fig. 2A




Fig. 2B  
SUBSTITUTE SHEET (RULE 26)



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C12N 15/12, C07K 14/51, 14/71</b> <b>G01N 33/566</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 97/07135</b> <b>(43) International Publication Date:</b> 27 February 1997 (27.02.97)
<p><b>(21) International Application Number:</b>      PCT/US96/13163</p> <p><b>(22) International Filing Date:</b>      13 August 1996 (13.08.96)</p> <p><b>(30) Priority Data:</b>              60/002,313      14 August 1995 (14.08.95)      US</p> <p><b>(71) Applicants:</b> CREATIVE BIOMOLECULES, INC. [US/US];              45 South Street, Hopkinton, MA 01748 (US). THE LUD-              WIG INSTITUTE FOR CANCER RESEARCH [US/US];              1345 Avenue of the Americas, New York, NY 10105 (US).</p> <p><b>(72) Inventors:</b> ICHUO, Hidenori; The Cancer Institute, Tokyo,              Japanese Foundation for Cancer Research, Dept. of Bio-              chemistry, 1-37-1 Kami-ikebukuro, Toshima-ku, Tokyo 170              (JP). NISHITOH, Hideki; The Cancer Institute, Tokyo,              Japanese Foundation for Cancer Research, Dept. of Bio-              chemistry, 1-37-1 Kami-ikebukuro, Toshima-ku, Tokyo 170              (JP). SAMPATH, Kuber, T.; 6 Spring Street, Medway, MA              02053 (US).</p> <p><b>(74) Agent:</b> GREENHALGH, Duncan, A.; Testa, Hurwitz &amp;              Thibault, L.L.P., High Street Tower, 125 High Street,              Boston, MA 02110 (US).</p>		<p><b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY,              CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU,              IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU,              LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO,              RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ,              VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian              patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European              patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT,              LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI,              CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b>  <i>With international search report.</i>  <i>Before the expiration of the time limit for amending the</i>  <i>claims and to be republished in the event of the receipt of</i>  <i>amendments.</i></p> <p><b>(88) Date of publication of the international search report:</b>              22 May 1997 (22.05.97)</p>
<p><b>(54) Title:</b> BINDING OF OSTEOGENIC PROTEIN-1 (OP-1) AND ANALOGS THEREOF TO THE CELL SURFACE RECEPTOR          ALK-1 AND ANALOGS THEREOF</p>		
<p><b>(57) Abstract</b></p> <p>Disclosed are (1) nucleic acid sequences, amino acid sequences, homologies, structural features, and various other data characterizing morphogen cell surface receptors, particularly OP-1-binding cell surface receptors; (2) methods for producing receptor proteins, including fragments thereof, using recombinant DNA technology; (3) methods for identifying novel morphogen receptors and their encoding DNAs; (4) methods for identifying compounds capable of modulating endogenous morphogen receptor levels; and (5) methods and compositions for identifying and producing morphogen analogs useful in the design of morphogen agonists and antagonists for therapeutic, diagnostic, and experimental uses.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

# INTERNATIONAL SEARCH REPORT

International Application No  
PC/US 96/13163

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/12 C07K14/51 C07K14/71 G01N33/566

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 07073 A (CREATIVE BIOMOLECULES) 30 April 1992 p. 7, last paragraph; p. 17., l. 29-30 - p. 18, l. 1-22; p. 20, l. 15-32; Figures 1 and 2	1-5
X	--- THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 25, 24 June 1994, pages 16985-16988, XP000611465 TEN DIJKE, P. ET AL.: "Identification of type I receptors for osteogenic protein-1 and bone morphogenetic protein-4" "Results" --- -/-	6-8, 10-14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*A\* document member of the same patent family

Date of the actual completion of the international search

24 January 1997

Date of mailing of the international search report

18. 04. 97

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl.  
Fax (+ 31-70) 340-3016

Authorized officer

Alt, G

## INTERNATIONAL SEARCH REPORT

Inter Application No

PC1/US 96/13163

## C.(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCIENCE, vol. 264, 1 April 1994, pages 101-104, XP000561968 TEN DIJKE, P. ET AL.: "Characterization of type I receptors for transforming growth factor-beta and activin" page 102, middle and right-hand column ---	6-8, 10-12
X	CELL, vol. 75, 19 November 1993, pages 671-680, XP000611962 ATTISANO, L. ET AL.: "Identification of human activin and TGF-beta type I receptors that form heteromeric kinase complexes with type II receptors" page 672, right-hand column - page 674, left-hand column, second paragraph ---	6-8, 10-12
A	ONCOGENE, vol. 8, October 1993, pages 2879-2887, XP000611724 TEN DIJKE, P. ET AL.: "Activin receptor-like kinases: a novel subclass of cell-surface receptors with predicted serine/threonine kinase activity" Figures 1 and 3 ---	1-14,17, 19,20
A	THE JOURNAL OF CELLULAR BIOLOGY, vol. 130, July 1995, pages 217-226, XP000611731 YAMASHITA, H. ET AL.: "Osteogenic protein-1 binds to activin type II receptors and induces certain activin-like effects" the whole document ---	1-14,17, 19,20
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 92, August 1995, pages 7632-7636, XP002021530 ROSENZWEIG, B.L. ET AL.: "Cloning and characterization of a human type II receptor for bone morphogenetic proteins" page 7634, left-hand column - page 7636 ---	1-14,17, 19,20
P,X	WO 95 30003 A (CREATIVE BIOMOLECULES) 9 November 1995 pages 6-16; Example 8 ---	1-14,17, 19,20
A	WO 94 06399 A (CREATIVE BIOMOLECULES) 31 March 1994 pages 11-17; pages 38-44 -----	1-14,17, 19,20

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/13163

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

- claims 1-14, 17, 19, 20
- claims 15, 15
- claim 18

\* see continuation-sheet PCT/ISA/210 \*

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-14, 17, 19, 20

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 96/ 13163

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

- claims 1-14,17,19,20: OP-1 analogs and methods of their application and for their identification and production
- claims 15,16 : A binding partner having specific binding affinity for an epitope on a ligand receptor complex but not to the uncomplexed OP-1 protein or analog
- claim 18 : A method for antagonising OP-1 binding by providing a protein having the ALK1 amino acid sequence or analogs thereof

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/US 96/13163

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9207073 A	30-04-92	AU 663689 B	19-10-95
		AU 8900091 A	20-05-92
		CA 2094027 A	19-04-92
		EP 0643767 A	22-03-95
		JP 7500487 T	19-01-95
		US 5468845 A	21-11-95
		US 5344654 A	06-09-94
		US 5354557 A	11-10-94
-----			
WO 9530003 A	09-11-95	AU 2466295 A	29-11-95
		CA 2187902 A	09-11-95
		EP 0756628 A	05-02-97
-----			
WO 9406399 A	31-03-94	AU 4795193 A	03-03-94
		AU 4797193 A	03-03-94
		AU 4995593 A	03-03-94
		AU 5129293 A	12-04-94
		AU 5129393 A	12-04-94
		AU 5162393 A	12-04-94
		AU 5290893 A	12-04-94
		AU 5590094 A	24-05-94
		CA 2141554 A	17-02-94
		CA 2141555 A	17-02-94
		CA 2141556 A	17-02-94
		CA 2147598 A	11-05-94
		EP 0652953 A	17-05-95
		EP 0653942 A	24-05-95
		EP 0661933 A	12-07-95
		EP 0665739 A	09-08-95
		EP 0661987 A	12-07-95
		EP 0680334 A	08-11-95
		EP 0672064 A	20-09-95
		JP 7509611 T	26-10-95
		JP 7509720 T	26-10-95
		JP 7509721 T	26-10-95
		JP 8501779 T	27-02-96
		JP 8501558 T	20-02-96
		JP 8501315 T	13-02-96
		JP 8503198 T	09-04-96
		WO 9403600 A	17-02-94



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/US 96/13163

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9406399 A		WO 9403075 A	17-02-94
		WO 9403200 A	17-02-94
		WO 9406447 A	31-03-94
		WO 9406449 A	31-03-94
		WO 9406420 A	31-03-94
		WO 9410203 A	11-05-94
		AU 670334 B	11-07-96
-----			